

THE CROSS COVARIOGRAM OF A PAIR OF POLYGONS DETERMINES BOTH POLYGONS, WITH A FEW EXCEPTIONS

GABRIELE BIANCHI

ABSTRACT. The cross covariogram $g_{K,L}$ of two convex sets K and L in $\mathbb{R}^n$ is the function which associates to each $x \in \mathbb{R}^n$ the volume of $K \cap (L + x)$. Very recently Averkov and Bianchi [AB] have confirmed Matheron's conjecture on the covariogram problem, that asserts that any planar convex body K is determined by the knowledge of $g_{K,K}$. The problem of determining the sets from their covariogram is relevant in probability, in statistical shape recognition and in the determination of the atomic structure of a quasicrystal from X-ray diffraction images. We prove that when K and L are convex polygons (and also when K and L are planar convex cones) $g_{K,L}$ determines both K and L , up to a described family of exceptions. These results imply that, when K and L are in these classes, the information provided by the cross covariogram is so rich as to determine not only one unknown body, as required by Matheron's conjecture, but two bodies, with a few classified exceptions. These results are also used by Bianchi [Bia] to prove that any convex polytope P in $\mathbb{R}^3$ is determined by $g_{P,P}$.

1. INTRODUCTION

Let K and L be convex sets in $\mathbb{R}^n$, $n \geq 2$, and let λ_n stand for the n -dimensional Lebesgue measure. The *cross covariogram* $g_{K,L}$ of K and L is the function defined by

$$g_{K,L}(x) = \lambda_n(K \cap (L + x)),$$

where $x \in \mathbb{R}^n$ is such that $\lambda_n(K \cap (L + x))$ is finite. This function, introduced by Cabo and Janssen [CJ94], coincides with the convolution of the characteristic function $\mathbf{1}_K$ of K with the characteristic function $\mathbf{1}_{-L}$ of the reflection of L in the origin, that is,

$$(1.1) \quad g_{K,L} = \mathbf{1}_K * \mathbf{1}_{-L}.$$

The function $g_{K,K}$ was introduced by G. Matheron in his book [Mat75, Section 4.3] on random sets, is denoted by g_K and is called *covariogram* or *set covariance* of K . The covariogram g_K is clearly unchanged by a translation or a reflection of K . (The term *reflection* will always mean reflection in a point.) A *convex body* in $\mathbb{R}^n$ is a convex compact set with non-empty interior. In 1986 Matheron [Mat86, p. 20] asked the following question and conjectured a positive answer for the case $n = 2$.

Covariogram problem. *Does g_K determine a convex body K , among all convex bodies, up to translations and reflections?*

Even for $n = 2$ this conjecture has been completely settled only very recently, by Averkov and Bianchi [AB]. It is known that the covariogram problem is equivalent to any of the following problems (see [AB] for a detailed explanation).

Date: May 13, 2008.

2000 Mathematics Subject Classification. Primary 60D05; Secondary 52A22, 52A10, 52A38.

Key words and phrases. covariogram, cross covariogram, geometric tomography, quasicrystal, phase retrieval, set covariance, synisothesis.

- P1. Determine a convex body K by the knowledge, for each unit vector u in $\mathbb{R}^n$, of the distribution of the lengths of the chords of K parallel to u .
- P2. Determine a convex body K by the distribution of $X - Y$, where X and Y are independent random variables uniformly distributed over K .
- P3. Determine the characteristic function $\mathbf{1}_K$ of a convex body K from the modulus of its Fourier transform $\widehat{\mathbf{1}_K}$.

Chord-length distributions are of wide interest beyond mathematics and are common in *stereology, statistical shape recognition and image analysis*, where properties of an unknown body have to be inferred from chord length measurements; see Schmitt [Scm93], Cabo and Baddeley [CB03], Serra [Ser84] and Mazzolo, Roesslinger and Gille [MRG03]. Problem P2 was asked by Adler and Pyke [AP91] in 1991. The same authors [AP97] find the covariogram problem relevant also in the study of scanning Brownian processes and of the equivalence of measures induced by these processes for different base sets. Problem P3 is a special case of the *phase retrieval problem*, where $\mathbf{1}_K$ is replaced by a function with compact support. The phase retrieval problem has applications in *X-ray crystallography, optics, electron microscopy* and other areas, references to which may be found in [BSV00]. Very recently, Baake and Grimm [BG07] have proved that the covariogram problem is particularly relevant for finding the atomic structure of a *quasicrystal* from its *X-ray* diffraction image. When a quasicrystal fits into the so-called *cut-and-project scheme*, the determination of its atomic structure S requires the knowledge of an unknown “window” W , which in many important cases is a convex body. The covariogram problem enters at this point, since g_W can be obtained from the diffraction image of S .

We have already mentioned that the covariogram problem has a positive answer in the plane. In higher dimensions a complete answer is known only when K is a convex polytope. Bianchi [Bia05] proved that in $\mathbb{R}^n$, for every $n \geq 4$, there are pairs of convex polytopes with equal covariograms which are not translations or reflections of each other. On the other hand, the answer to the covariogram problem for a three-dimensional convex polytope is positive, as proved by Bianchi [Bia].

Cabo and Janssen [CJ94] prove that $g_{C,-C}$ determines every regular (equal to the closure of its interior) compact set C in $\mathbb{R}^n$, $n \geq 2$. This result clearly implies that the covariogram determines each centrally symmetric regular compact set in $\mathbb{R}^n$. In general, the convexity assumption in the covariogram problem is needed, since there exist examples of non-convex polyominoes which are neither translations nor reflections of each other and have equal covariograms; see Gardner, Gronchi and Zong [GGZ05]. However, a planar convex body is determined by its covariogram in a class that is much larger than that of convex bodies; see Benassi, Bianchi and D’Ercole [BBD].

When K is a planar convex body, the information provided by g_K seems to be richer than is necessary to determine K . For instance, for a convex body K whose boundary is C^2 regular and has non-zero curvature, Averkov and Bianchi [AB07] indicate some subsets of the support of g_K , with arbitrarily small Lebesgue measure, such that g_K restricted to those subsets identifies K . In this paper we investigate this richness of information from a different point of view, trying to understand which information $g_{K,L}$ carries about the *two* convex sets K and L . We are able to provide a complete answer when K and L are convex polygons, and also when they are planar convex cones. The obtained results imply that the information provided by the cross covariogram, when K and L are in these classes, is so rich as to determine not only one unknown body, as required by Matheron’s conjecture, but two bodies, with a few classified exceptions.

In order to generalise the covariogram problem to the case of the cross covariogram, we observe that the translation of K and L by the same vector, and the substitution of K with $-L$ and of L with $-K$, leave $g_{K,L}$ unchanged. Let K , L , K' and L' be convex sets in $\mathbb{R}^2$. We call (K, L) and (K', L') *trivial associates* when one pair is obtained by the other one via a combination of the two operations above, that is, when either $(K, L) = (K' + x, L' + x)$ or $(K, L) = (-L' + x, -K' + x)$, for some $x \in \mathbb{R}^2$.

Cross covariogram problem. *Does $g_{K,L}$ determine the pair of closed convex sets (K, L) , among all pairs of closed convex sets, up to trivial associates?*

Assume K and L convex polygons. In this case the answer to the cross covariogram problem is negative as Examples 4.1 and 5.2 show (see Figures 3 and 4). For each choice of some real parameters there exist four pairs of parallelograms $(\mathcal{K}_1, \mathcal{L}_1), \dots, (\mathcal{K}_4, \mathcal{L}_4)$ such that, for $i = 1, 3$, $g_{\mathcal{K}_i, \mathcal{L}_i} = g_{\mathcal{K}_{i+1}, \mathcal{L}_{i+1}}$ but $(\mathcal{K}_i, \mathcal{L}_i)$ is not a trivial associate of $(\mathcal{K}_{i+1}, \mathcal{L}_{i+1})$. Theorem 1.1 proves that, up to an affine transformation, the previous counterexamples are the only ones.

Theorem 1.1. *Let K and L be convex polygons and K' and L' be planar closed convex sets with $g_{K,L} = g_{K',L'}$. Assume that there exist no affine transformation T and no different indices i, j , with either $i, j \in \{1, 2\}$ or $i, j \in \{3, 4\}$, such that (TK, TL) and (TK', TL') are trivial associates of $(\mathcal{K}_i, \mathcal{L}_i)$ and of $(\mathcal{K}_j, \mathcal{L}_j)$, respectively. Then (K, L) is a trivial associate of (K', L') .*

Theorem 1.1 has a probabilistic interpretation in terms of a generalisation of Problem P2. It implies that the distribution of the difference $X - Y$ of two independent random variables X and Y , with X uniformly distributed over a convex polygon K and Y uniformly distributed over a convex polygon L , together with $\lambda_2(K) \lambda_2(L)$, determines both K and L , up to some inherent ambiguities, with a few exceptions. This result is a consequence of Theorem 1.1 because the probability distribution of $X - Y$ is $g_{K,L}/(\lambda_2(K) \lambda_2(L))$, by (1.1).

It would be interesting to understand if a result similar to Theorem 1.1 holds for other classes of planar convex bodies, for instance for bodies with C^2 boundary.

Another motivation for Theorem 1.1 comes from the proof of the positive answer to the covariogram problem for three-dimensional convex polytopes mentioned above, of which Theorem 1.1 constitutes a crucial step. The two problems are connected because when K and L are parallel antipodal facets of a three-dimensional convex polytope P , g_P provides $g_{K,L}$, and Theorem 1.1 helps to determine those pairs of facets; see [Bia] for the details.

The previous theorem has something to say also regarding the symmetries of $g_{K,L}$. Let F and G be convex bodies in $\mathbb{R}^n$. It is evident that $g_F(x) = g_F(-x)$ for every $x \in \mathbb{R}^n$, but the cross covariogram is not always an even function. (In general, one only has $g_{F,G}(-x) = g_{G,F}(x) = g_{-F,-G}(x)$.)

Corollary 1.2. *Let K and L be convex polygons. Then there exists $z \in \mathbb{R}^2$ such that $g_{K,L}(z+x) = g_{K,L}(z-x)$ for every $x \in \mathbb{R}^2$ if and only if either $K = L + z$ or both K and $L + z$ are centrally symmetric with equal center.*

Mani-Levitska [Man01] saw the study of the cross covariogram problem for pairs of polyhedral convex cones in $\mathbb{R}^3$ as a step towards the solution of the covariogram problem for polytopes in $\mathbb{R}^3$, and indeed [Bia] contains some results in this direction. These results for cones in $\mathbb{R}^3$ are not exhaustive, but the cross covariogram problem for planar convex cones can be completely understood. Let A , A' , B and B' be convex cones in $\mathbb{R}^2$ with apex O . Assume $\text{int } A \cap \text{int } B = \emptyset$, because otherwise $g_{A,B}$ is nowhere finite. Since the cones have apex O , (A, B) and (A', B') are trivial associates if and only if $\{A, -B\} = \{A', -B'\}$. Example 3.1 (see Figure 1) presents two different pairs of convex cones $(\mathcal{A}_1, \mathcal{B}_1)$ and $(\mathcal{A}_2, \mathcal{B}_2)$ with equal cross

covariogram and Theorem 1.3 proves that, up to affine transformations, these are the only counterexamples.

Theorem 1.3. *Let A , B , A' and B' be pointed closed convex cones in $\mathbb{R}^2$ with non-empty interior and apex the origin O , such that $\text{int } A \cap \text{int } B = \emptyset$. The identity $g_{A,B} = g_{A',B'}$ holds true if and only if one of the following alternatives occurs:*

- (i) $\{A, -B\} = \{A', -B'\}$;
 - (ii) *there exist a linear transformation $\mathcal{T}$ and $i, j \in \{1, 2\}$, $i \neq j$, such that*
- $$(1.2) \quad \{\mathcal{T}A, -\mathcal{T}B\} = \{\mathcal{A}_i, -\mathcal{B}_i\} \quad \text{and} \quad \{\mathcal{T}A', -\mathcal{T}B'\} = \{\mathcal{A}_j, -\mathcal{B}_j\}.$$

A crucial notion in the proof of Theorem 1.1 is that of *synisothetic* pairs of polytopes, introduced by [Bia] and explained in Section 2. In Section 4 we prove that, up to affine transformations, $(\mathcal{K}_1, \mathcal{L}_1)$ and $(\mathcal{K}_2, \mathcal{L}_2)$ are the only pairs of convex polygons with equal cross covariogram which are not synisothetic. To establish this result we use also Theorem 1.3, whose proof is contained in Section 3. The proofs of Theorem 1.1 and of Corollary 1.2 are contained in Section 6. We conclude by mentioning that Lemma 5.1 is a technical result which may be of interest by itself.

2. DEFINITIONS, NOTATIONS AND PRELIMINARIES

As usual, S^{n-1} denotes the unit sphere in $\mathbb{R}^n$, centred at the origin O . For $x, y \in \mathbb{R}^n$, $\|x\|$ is the Euclidean norm of x and $x \cdot y$ denotes scalar product. For $\delta > 0$, $B(x, \delta)$ denotes the open ball in $\mathbb{R}^n$ centred at x and with radius δ . For $\theta \in [0, 2\pi]$ we write $u(\theta) = (\cos \theta, \sin \theta) \in S^1$.

If $A \subset \mathbb{R}^n$ we denote by $\text{int } A$, $\text{cl } A$, ∂A and $\text{conv } A$ the *interior*, *closure*, *boundary* and *convex hull* of A , respectively. The *symmetric difference* of the sets A and B is defined by $A \Delta B = (A \setminus B) \cup (B \setminus A)$. The *Minkowski sum* of A and B is

$$A + B = \{x + y : x \in A, y \in B\}.$$

We write λ_k for k -dimensional Lebesgue measure in $\mathbb{R}^n$, where $k = 1, \dots, n$, and where we identify λ_k with k -dimensional Hausdorff measure.

A *convex body* $K \subset \mathbb{R}^n$ is a compact convex set with non-empty interior. The symbols $\text{relbd } K$ and $\text{relint } K$ indicate respectively the *relative boundary* and the *relative interior* of K . The *difference body* of K is defined by $DK = K + (-K)$. The *support function* of K is defined, for $x \in \mathbb{R}^n$, by $h_K(x) = \sup \{x \cdot y : y \in K\}$. Given $x, y \in \mathbb{R}^n$, we write $[x, y]$ for the line segment with endpoints x and y . When K is a planar convex body and $a, b \in \partial K$, the symbol $(a, b)_{\partial K}$ denotes the set of points in ∂K which strictly follow a and strictly precede b in counterclockwise order on ∂K , and $[a, b]_{\partial K}$ denotes $(a, b)_{\partial K} \cup \{a, b\}$. Given an arc $\Omega \subset \partial K$ with $\text{cl } \Omega = [a, b]_{\partial K}$, we call a the *lower endpoint* of Ω , b its *upper endpoint*, and, with a small abuse of notation, we will call $(a, b)_{\partial K}$ the *relative interior* of Ω . Given $u, v \in S^1$, $v \geq u$ means $v \in (u, -u)_{S^1} \cup \{u\}$, while $v > u$ means $v \in (u, -u)_{S^1} \cup \{-u\}$.

If F is a face of a convex polytope P in $\mathbb{R}^n$, the *normal cone* of P at F is denoted by $N(P, F)$ and is the set of all outer normal vectors to P at x , where $x \in \text{relint } F$, together with O . The *support cone* of P at F is the set

$$\text{cone}(P, F) = \{\mu(y - x) : y \in P, \mu \geq 0\},$$

where $x \in \text{relint } F$. Neither definitions depend on the choice of x . If $u \in S^{n-1}$, the *exposed face* of P in direction u is

$$P_u = \{x \in P : x \cdot u = h_P(u)\}.$$

It is the unique proper face of P such that the relative interior of its normal cone contains u . [Sch93, Th. 1.7.5(c)] proves that, when K and L are convex polygons and $u \in S^1$, we have

$$(2.1) \quad (K + (-L))_u = K_u + (-L)_u.$$




FIGURE 1. Up to affine transformations, these are the only different pairs of convex cones with equal cross covariogram.

In this paper the term *cone* always means cone with apex O . A convex cone is *pointed* if its apex is a vertex. Let (ρ, θ) denote polar coordinates in $\mathbb{R}^2$ and let $\alpha, \beta \in [0, 2\pi]$, with $\alpha < \beta$. For brevity we write $\{\theta = \alpha\}$ for the ray $\{(\rho, \theta) : \theta = \alpha\}$ and $\{\alpha \leq \theta \leq \beta\}$ for the cone $\{(\rho, \theta) : \theta \in [\alpha, \beta]\}$.

Let $\text{supp } f$ denote the support of the function f . Let K and L be convex polygons and let A and B be closed convex cones in $\mathbb{R}^2$ with $\text{int } A \cap \text{int } B = \emptyset$. It is easy to prove that

$$(2.2) \quad \text{supp } g_{K,L} = K + (-L) \quad \text{and} \quad \text{supp } g_{A,B} = A + (-B) = \text{conv}(A \cup (-B)).$$

It can be proved, by using the Minkowski inequality as in [Sch93, p. 410-411], that $g_{K,L}^{1/2}$ is concave on its support.

Synisothesis. Let P and Q be convex polytopes in $\mathbb{R}^n$, let F be a proper face of P , and let G be a proper face of Q . We say that F and G are *isothetic* if G is a translate of F and

$$\text{cone}(P, F) = \text{cone}(Q, G).$$

Given convex polytopes P_1, P_2, Q_1 and Q_2 in $\mathbb{R}^n$ we say that (P_1, P_2) and (Q_1, Q_2) are *synisothetic* if given any proper face F of P_1 or of P_2 there is a proper face G of Q_1 or of Q_2 (and conversely) such that F and G are isothetic.

The notion of synisothesis will play a central role in this paper. Let K, K', L and L' be convex polygons. It is convenient for later use to express the synisothesis of $(K, -L)$ and $(K', -L')$ in two equivalent ways. It is clear that $(K, -L)$ and $(K', -L')$ are synisothetic if and only if, for each $u \in S^1$, one of the following properties hold:

$$(2.3) \quad K_u \text{ is isothetic to } K'_u \text{ and } (-L)_u \text{ is isothetic to } (-L')_u;$$

$$(2.4) \quad K_u \text{ is isothetic to } (-L')_u \text{ and } (-L)_u \text{ is isothetic to } K'_u.$$

Let $u \in S^1$ and F and F' be convex polygons. Observe that F_u is isothetic to F'_u if and only if either both are edges of equal length or both are vertices with equal support cones. Thus, $(K, -L)$ and $(K', -L')$ are synisothetic if and only if the following equalities hold for each $u \in S^1$:

$$(2.5) \quad \{\lambda_1(K_u), \lambda_1((-L)_u)\} = \{\lambda_1(K'_u), \lambda_1((-L')_u)\};$$

$$(2.6) \quad \{\text{cone}(K, K_u), \text{cone}(-L, (-L)_u)\} = \{\text{cone}(K', K'_u), \text{cone}(-L', (-L')_u)\}.$$

3. THE CROSS COVARIOGRAM PROBLEM FOR PLANAR CONVEX CONES

Let us introduce the counterexample to the cross covariogram problem for cones.

Example 3.1. Let $\mathcal{A}_1 = \{0 \leq \theta \leq 3\pi/4\}$, $\mathcal{B}_1 = -\{\pi/4 \leq \theta \leq \pi/2\}$, $\mathcal{A}_2 = \{0 \leq \theta \leq \pi/4\}$ and $\mathcal{B}_2 = -\{\pi/2 \leq \theta \leq 3\pi/4\}$; see Fig. 1. Clearly $\{\mathcal{A}_1, -\mathcal{B}_1\} \neq \{\mathcal{A}_2, -\mathcal{B}_2\}$. Ele-

mentary calculations prove $g_{\mathcal{A}_1, \mathcal{B}_1} = g_{\mathcal{A}_2, \mathcal{B}_2}$. Indeed, if $x = (x_1, x_2)$ in Cartesian coordinates, then

$$g_{\mathcal{A}_1, \mathcal{B}_1}(x) = g_{\mathcal{A}_2, \mathcal{B}_2}(x) = \begin{cases} x_2^2/2 & \text{if } x \in \{0 \leq \theta \leq \pi/4\}; \\ (x_2^2 - x_1^2 + 2x_1x_2)/4 & \text{if } x \in \{\pi/4 \leq \theta \leq \pi/2\}; \\ (x_1 + x_2)^2/4 & \text{if } x \in \{\pi/2 \leq \theta \leq 3\pi/4\}; \\ 0 & \text{if } x \notin \{0 \leq \theta \leq 3\pi/4\}. \end{cases}$$

These properties are preserved by any non-singular affine transformation $\mathcal{T}$, since

$$g_{\mathcal{T}\mathcal{A}_1, \mathcal{T}\mathcal{B}_1}(x) = |\det \mathcal{T}| g_{\mathcal{A}_1, \mathcal{B}_1}(\mathcal{T}^{-1}x) = |\det \mathcal{T}| g_{\mathcal{A}_2, \mathcal{B}_2}(\mathcal{T}^{-1}x) = g_{\mathcal{T}\mathcal{A}_2, \mathcal{T}\mathcal{B}_2}(x).$$

Lemma 3.2. *Let A and B be pointed closed convex cones in $\mathbb{R}^2$ with non-empty interior satisfying $A, -B \subset \{0 \leq \theta \leq \pi\}$. The set $S^2(A, B) = \text{cl}\{x \in \mathbb{R}^2 : g_{A, B} \text{ is not } C^2 \text{ at } x\}$ coincides with $\partial A \cup (-\partial B)$.*

Proof. Let $W \subset \mathbb{R}^2 \setminus (\partial A \cup (-\partial B))$ be an open connected set. When $x \in W$ the vertex x of $B + x$ does not belong to ∂A , and the vertex O of A does not belong to $\partial B + x$. Thus the combinatorial structure of $\partial(A \cap (B + x))$ is constant in W . Since the vertices of $A \cap (B + x)$ are smooth functions of x , for each $x \in W$, so is $g_{A, B}$. This proves $S^2(A, B) \subset \partial A \cup (-\partial B)$.

Let $B = \{\alpha \leq \theta \leq \beta\}$, for suitable $\alpha, \beta \in [\pi, 2\pi]$ with $\alpha - \beta \neq \pm\pi$. It is easy to prove that, when $x \notin \partial A$, we have

$$\begin{aligned} \frac{\partial^2 g_{A, B}}{\partial u(\alpha) \partial u(\beta)}(x) &= -|\sin(\alpha - \beta)| \frac{\partial}{\partial u(\alpha)} \lambda_1(A \cap (x + \{\theta = \beta\})) \\ &= -|\sin(\alpha - \beta)| 1_A(x), \end{aligned}$$

This formula proves $\partial A \subset S^2(A, B)$. A similar formula for the second order mixed derivative of $g_{A, B}$ in the directions of the edges of A proves $-\partial B \subset S^2(A, B)$. $\square$


Proof of Theorem 1.3. We have $\text{conv}(A \cup (-B)) = \text{conv}(A' \cup (-B')) = \text{supp } g_{A, B}$, by (2.2). Choose polar coordinates (ρ, θ) so that $\text{supp } g_{A, B} \subset \{0 \leq \theta \leq \pi\}$. Lemma 3.2 proves that $\partial A \cup (-\partial B)$ is determined by $g_{A, B}$. If $\partial A \cup (-\partial B)$ consists of two rays, then both A and $-B$ coincide with the convex cone bounded by those rays. Therefore $\{A, -B\}$ is determined by $g_{A, B}$ and $g_{A, B} = g_{A', B'}$ implies (i).

Assume that $\partial A \cup (-\partial B)$ consists of three rays. Let $0 \leq \theta_1 < \theta_2 < \theta_3 \leq \pi$ be the angles corresponding to these rays. Clearly exactly one among the rays that bound A and $-B$ coincides with $\partial A \cap (-\partial B)$. Elementary calculations show that, as $\varepsilon \rightarrow 0^+$,

$$g_{A, B}(u(\theta_1 + \varepsilon)) = \begin{cases} \varepsilon + o(\varepsilon) & \text{if } \{\theta = \theta_1\} = \partial A \cap (-\partial B); \\ o(\varepsilon) & \text{otherwise.} \end{cases}$$

An analogous formula, with $\{\theta = \theta_3\}$ substituting $\{\theta = \theta_1\}$, holds for $g_{A, B}(u(\theta_3 - \varepsilon))$. From the asymptotic behaviour of $g_{A, B}(u(\theta_1 + \varepsilon))$ and $g_{A, B}(u(\theta_3 - \varepsilon))$ it is thus possible to understand which of the three rays $\{\theta = \theta_1\}$, $\{\theta = \theta_2\}$ and $\{\theta = \theta_3\}$ coincides with $\partial A \cap (-\partial B)$. If, for instance, $\{\theta = \theta_2\} \subset \partial A \cap (-\partial B)$, then we necessarily have either $A = \{\theta_1 \leq \theta \leq \theta_2\}$ and $-B = \{\theta_2 \leq \theta \leq \theta_3\}$ or else $-B = \{\theta_1 \leq \theta \leq \theta_2\}$ and $A = \{\theta_2 \leq \theta \leq \theta_3\}$. Thus $\{A, -B\}$ is determined. Similar arguments prove that $\{A, -B\}$ is determined when $\partial A \cap (-\partial B)$ coincides with $\{\theta = \theta_1\}$ or with $\{\theta = \theta_3\}$. The equality $g_{A, B} = g_{A', B'}$ implies (i).

Assume that $\partial A \cup (-\partial B)$ consists of four rays, say $\{\theta = \theta_i\}$, $i = 1, \dots, 4$, with $0 \leq \theta_1 < \theta_2 < \theta_3 < \theta_4 \leq \pi$. Let P_2 be the parallelogram bounded by the rays $\{\theta = \theta_1\}$ and $\{\theta = \theta_4\}$ and by the lines which are parallel to these rays and contain $u(\theta_2)$ (see Fig. 2). Let $a_2 \neq O$ be the vertex of P_2 in $\{\theta = \theta_4\}$, $b_2 \neq O$ be the vertex

FIGURE 2. The set $A \cap (B + u(\theta_2))$ in Case 1 (left) and in Case 2 (right).

of P_2 in $\{\theta = \theta_1\}$ and let $c_2 = \{\theta = \theta_3\} \cap [u(\theta_2), a_2]$. There are three possible cases.

Case 1. $\{A, -B\} = \{\{\theta_1 \leq \theta \leq \theta_3\}, \{\theta_2 \leq \theta \leq \theta_4\}\}$.

It is easy to see that $A \cap (B + u(\theta_2))$ is equal to the triangle $\text{conv}\{O, u(\theta_2), b_2\}$ when $A = \{\theta_1 \leq \theta \leq \theta_3\}$ and $-B = \{\theta_2 \leq \theta \leq \theta_4\}$ (see Fig. 2(a)), and it is equal to the triangle $\text{conv}\{O, u(\theta_2), a_2\}$ when $A = \{\theta_2 \leq \theta \leq \theta_4\}$ and $-B = \{\theta_1 \leq \theta \leq \theta_3\}$. In each case we have

$$(3.1) \quad g_{A,B}(u(\theta_2)) = \frac{1}{2} \lambda_2(P_2).$$

Case 2. $\{A, -B\} = \{\{\theta_2 \leq \theta \leq \theta_3\}, \{\theta_1 \leq \theta \leq \theta_4\}\}$.

Suppose that $A = \{\theta_2 \leq \theta \leq \theta_3\}$ and $-B = \{\theta_1 \leq \theta \leq \theta_4\}$. The set $A \cap (B + u(\theta_2))$ is strictly contained in the triangle $T = \text{conv}(O, u(\theta_2), a_2)$ (see Fig. 2(b)). Moreover the ratio between $\lambda_2(A \cap (B + u(\theta_2)))$ and $\lambda_2(T)$ equals the ratio between $\|u(\theta_2) - c_2\|$ and $\|u(\theta_2) - a_2\|$. Thus

$$(3.2) \quad g_{A,B}(u(\theta_2)) = \frac{\|u(\theta_2) - c_2\|}{2\|u(\theta_2) - a_2\|} \lambda_2(P_2) < \frac{1}{2} \lambda_2(P_2).$$

The same formulas also hold when $-B = \{\theta_2 \leq \theta \leq \theta_3\}$ and $A = \{\theta_1 \leq \theta \leq \theta_4\}$.

Case 3. $\{A, -B\} = \{\{\theta_1 \leq \theta \leq \theta_2\}, \{\theta_3 \leq \theta \leq \theta_4\}\}$.

Arguments similar to those of Case 2 prove the following formula:

$$(3.3) \quad g_{A,B}(u(\theta_2)) = \frac{\|c_2 - a_2\|}{2\|u(\theta_2) - a_2\|} \lambda_2(P_2) < \frac{1}{2} \lambda_2(P_2).$$

The comparison of the values of $g_{A,B}(u(\theta_2))$ in (3.1), (3.2) and (3.3) distinguishes Case 1 from the others. Moreover, it distinguishes Case 2 from Case 3 except when c_2 divides the segment $[u(\theta_2), a_2]$ in two equal parts. Assume that this happens and let $\mathcal{T}$ be a non-singular linear transformation which maps the ray $\{\theta = \theta_i\}$ in $\{\theta = (i-1)\pi/4\}$, for $i = 1, 2, 4$. The assumption regarding c_2 easily implies $\mathcal{T}\{\theta = \theta_3\} = \{\theta = \pi/2\}$.

The same analysis, with the same θ_i , is also valid for A' and B' . If the same case applies to (A, B) and to (A', B') , then $\{A, -B\} = \{A', -B'\}$. The only possibility left is that there exists an affine transformation $\mathcal{T}$ such that $\mathcal{T}\{\theta = \theta_i\} = \{\theta = (i-1)\pi/4\}$, for $i = 1, \dots, 4$. Case 2 applies to (A, B) and Case 3 applies to (A', B') (or vice versa). If this happens, then Alternative (ii) in Theorem 1.3 holds. $\square$

Remark 3.3. Observe that $\text{int } \mathcal{A}_1 \cap \text{int } (-\mathcal{B}_1) \neq \emptyset$ and $\text{int } \mathcal{A}_2 \cap \text{int } (-\mathcal{B}_2) = \emptyset$. Thus, if $\text{int } A \cap \text{int } (-B)$ and $\text{int } A' \cap \text{int } (-B')$ are both empty or both non-empty, then Theorem 1.3 implies $\{A, -B\} = \{A', -B'\}$.




FIGURE 3. Up to affine transformations, $(\mathcal{K}_1, \mathcal{L}_1)$ and $(\mathcal{K}_2, \mathcal{L}_2)$ are the only pairs of convex polygons with equal cross covariogram which are not synisothetic.

Remark 3.4. In the previous proof it is clear that the linear map $\mathcal{T}$ in Theorem 1.3 preserves the order of the rays, that is, the ray $\mathcal{T}^{-1}\{\theta = i\pi/4\}$ follows in counterclockwise order the ray $\mathcal{T}^{-1}\{\theta = (i-1)\pi/4\}$, for $i = 1, 2, 3$.

4. CROSS COVARIOGRAM AND SYNISOTHESES

The next example is due to this author and R. J. Gardner.

Example 4.1. Let α, β, γ and δ be positive real numbers, $I_1 = [(-1, 0), (1, 0)]$, $I_2 = 1/\sqrt{2} [(-1, -1), (1, 1)]$, $I_3 = [(0, -1), (0, 1)]$ and $I_4 = 1/\sqrt{2} [(1, -1), (-1, 1)]$. We define four parallelograms as follows:

$$\mathcal{K}_1 = \alpha I_1 + \beta I_2; \quad \mathcal{L}_1 = \gamma I_3 + \delta I_4 + y; \quad \mathcal{K}_2 = \alpha I_1 + \delta I_4 \quad \text{and} \quad \mathcal{L}_2 = \beta I_2 + \gamma I_3 + y.$$

See Fig. 3. The pairs $(\mathcal{K}_1, -\mathcal{L}_1)$ and $(\mathcal{K}_2, -\mathcal{L}_2)$ are not synisothetic (no vertex in the second pair has a support cone equal to the support cone of the top vertex of $\mathcal{L}_1$ or to its reflection). Moreover $g_{\mathcal{K}_1, \mathcal{L}_1} = g_{\mathcal{K}_2, \mathcal{L}_2}$. To prove it, let X_i and Y_i be independent random variables uniformly distributed over $\mathcal{K}_i$ and $\mathcal{L}_i$, for $i = 1, 2$, and let $Z_1, \dots, Z_4$ be independent random variables uniformly distributed over $I_1, \dots, I_4$, respectively. Then $X_1 = \alpha Z_1 + \beta Z_2$, $Y_1 = \gamma Z_3 + \delta Z_4 + y$, $X_2 = \alpha Z_1 + \delta Z_4$ and $Y_2 = \beta Z_2 + \gamma Z_3 + y$. Moreover we have

$$(4.1) \quad X_1 - Y_1 = X_2 - Y_2,$$

because $Z_2 = -Z_2$ and $Z_4 = -Z_4$. The distribution of probability of $X_i - Y_i$ is $1_{\mathcal{K}_i} * 1_{-\mathcal{L}_i} / (\lambda_2(\mathcal{K}_i) \lambda_2(\mathcal{L}_i))$. Observe that $\lambda_2(\mathcal{K}_1) \lambda_2(\mathcal{L}_1) = \alpha \beta \gamma \delta / 2 = \lambda_2(\mathcal{K}_2) \lambda_2(\mathcal{L}_2)$. Therefore (4.1) and (1.1) imply

$$g_{\mathcal{K}_1, \mathcal{L}_1} = 1_{\mathcal{K}_1} * 1_{-\mathcal{L}_1} = 1_{\mathcal{K}_2} * 1_{-\mathcal{L}_2} = g_{\mathcal{K}_2, \mathcal{L}_2}.$$

Proposition 4.2. Let K and L be convex polygons, K' and L' be planar closed convex sets with $g_{K, L} = g_{K', L'}$. Then K' and L' are polygons. Assume, moreover, that there is no affine transformation $\mathcal{T}$ and no different indices $i, j \in \{1, 2\}$ such that $(\mathcal{T}K, \mathcal{T}L)$ and $(\mathcal{T}K', \mathcal{T}L')$ are trivial associates of $(\mathcal{K}_i, \mathcal{L}_i)$ and $(\mathcal{K}_j, \mathcal{L}_j)$, respectively. Then $(K, -L)$ and $(K', -L')$ are synisothetic.

This proof is divided in three steps and occupies all the rest of the section. We recall that $(K, -L)$ and $(K', -L')$ are synisothetic if and only if for each $u \in S^1$ both (2.5) and (2.6) hold. We will prove that (2.5) holds for each u , while (2.6) fails for some u exactly when $(\mathcal{T}K, \mathcal{T}L)$ and $(\mathcal{T}K', \mathcal{T}L')$ are trivial associates of $(\mathcal{K}_i, \mathcal{L}_i)$ and $(\mathcal{K}_j, \mathcal{L}_j)$, respectively.

Observe that the set $K' - L'$ is a polygon, because $K' - L' = K - L$ by (2.2). This may happen only if K' and L' are polygons. The function $g_{K, L}$ determines

its support $K - L$ and, for all $u \in S^1$, it determines also $\lambda_1(K_u) + \lambda_1(L_{-u})$, since (2.1) implies

$$(4.2) \quad \lambda_1((K - L)_u) = \lambda_1(K_u) + \lambda_1(L_{-u}).$$

Claim 4.2.1. *The function $g_{K,L}$ determines $\{\lambda_1(K_u), \lambda_1(L_{-u})\}$ for each $u \in S^1$.*

Proof. If $(K - L)_u$ is a vertex, then both K_u and L_{-u} are vertices, by (4.2). In this case $\{\lambda_1(K_u), \lambda_1(L_{-u})\} = \{0\}$. Assume that

$$(K - L)_u \text{ is an edge.}$$

In this case at least one among K_u and L_{-u} is an edge. Let x be the midpoint of $(K - L)_u$. For sufficiently small $\varepsilon > 0$, let $x_\varepsilon \in [O, x]$ be the point at distance ε from $(K - L)_u$. Then it is easy to see that, as $\varepsilon \rightarrow 0^+$, we have

$$(4.3) \quad g_{K,L}(x_\varepsilon) = \min\{\lambda_1(K_u), \lambda_1(L_{-u})\}\varepsilon + o(\varepsilon).$$

(This corresponds to translating L so that the translated midpoint of L_{-u} is close to the midpoint of K_u .)

If $g_{K,L}(x_\varepsilon) = o(\varepsilon)$, then $\min\{\lambda_1(K_u), \lambda_1(L_{-u})\} = 0$, that is, either K_u is an edge and L_{-u} is a vertex, or vice versa. If, for instance, L_{-u} is a vertex, then the length of K_u is determined, since it equals the length of $(K - L)_u$.

If $g_{K,L}(x_\varepsilon) \geq \alpha\varepsilon$, for some constant $\alpha > 0$, then both K_u and L_{-u} are edges. From (4.3) we obtain the minimum of the lengths of these edges. Since the sum of these lengths is determined by (4.2), the pair $\{\lambda_1(K_u), \lambda_1(L_{-u})\}$ is known. $\square$

Let $z_0, \dots, z_n, w_0, \dots, w_t, z'_0, \dots, z'_{n'}, w'_0, \dots, w'_{t'}$ and $q_0, \dots, q_p$ denote respectively the vertices of K , of L , of K' , of L' and of $K - L = K' - L'$ in counterclockwise order on the respective boundaries. For each j , let $A_j = \text{cone}(K, z_j)$, $B_j = \text{cone}(L, w_j)$, $A'_j = \text{cone}(K', z'_j)$ and $B'_j = \text{cone}(L', w'_j)$. Each vertex of $K - L$ is the difference of a vertex of K and of one of L , and also of a vertex of K' and of one of L' . Assume that $q_m = z_s - w_l = z'_h - w'_k$. In a neighbourhood of z_s , K equals $A_s + z_s$, while in a neighbourhood of w_l , L equals $B_l + w_l$. If x belongs to a small neighbourhood of q_m , then we have

$$K \cap (L + x) = (A_s + z_s) \cap (B_l + w_l + x)$$

and this set is a translate of $A_s \cap (B_l + x - q_m)$. The function g_{A_s, B_l} is thus determined, in a neighbourhood of O , by $g_{K,L}$. Since g_{A_s, B_l} is homogeneous of degree 2 it is determined on its entire domain. Similar considerations apply to (K', L') and imply $g_{A_s, B_l} = g_{A'_h, B'_k}$.

We say that the vertex $q_m = z_s - w_l = z'_h - w'_k$ is *ambiguous* if $\{A_s, -B_l\} \neq \{A'_h, -B'_k\}$.

It is elementary to check that when $[z_s, z_{s+1}]$ and $[w_l, w_{l+1}]$ are parallel we have

$$(4.4) \quad [q_m, q_{m+1}] = [z_s, z_{s+1}] - [w_l, w_{l+1}],$$

while when they are not parallel we have either

$$(4.5) \quad [q_m, q_{m+1}] = [z_s, z_{s+1}] - w_l \text{ or}$$

$$(4.6) \quad [q_m, q_{m+1}] = z_s - [w_l, w_{l+1}].$$

Claim 4.2.2. *Let $q_m = z_s - w_l$ and assume that q_m is ambiguous. Then q_{m+1} is ambiguous too and (4.4) does not hold. Moreover, when (4.5) holds, $[z_{s-1}, z_s]$ and $[z_{s+1}, z_{s+2}]$ are parallel, while when (4.6) holds, $[w_{l-1}, w_l]$ and $[w_{l+1}, w_{l+2}]$ are parallel.*

Proof. Choose polar coordinates (ρ, θ) so that $[q_m, q_{m+1}]$ is parallel to $\{\theta = 0\}$ and A_s and $-B_l$ are contained in $\{0 \leq \theta \leq \pi\}$. Since $\text{conv}(A_s \cup (-B_l)) = \text{conv}(A'_h \cup (-B'_k))$, by (2.2), also A'_h and $-B'_k$ are contained in $\{0 \leq \theta \leq \pi\}$. Theorem 1.3

proves that there exist a linear transformation $\mathcal{T}$ and $i, j \in \{1, 2\}$, with $i \neq j$, such that Alternative (ii) of Theorem 1.3 occurs, with $A = A_s$, $B = B_l$, $A' = A'_h$ and $B' = B'_k$. Since no edge of $\mathcal{A}_i$ is parallel to an edge of $-\mathcal{B}_i$, for each choice of i , (1.2) implies that no edge of K adjacent to z_s is parallel to an edge of L adjacent to w_l . This rules out (4.4). Analogous arguments rule out $[q_m, q_{m+1}] = [z'_h, z'_{h+1}] - [w'_k, w'_{k+1}]$. The latter implies that $[q_m, q_{m+1}]$ equals either $[z'_h, z'_{h+1}] - w'_k$ or $z'_h - [w'_k, w'_{k+1}]$.

Assume (4.5) and $[q_m, q_{m+1}] = [z'_h, z'_{h+1}] - w'_k$. This clearly implies

$$(4.7) \quad A_s = \{0 \leq \theta \leq \alpha\}, \quad A'_h = \{0 \leq \theta \leq \alpha'\},$$

$$(4.8) \quad A_{s+1} = \{\beta \leq \theta \leq \pi\} \quad \text{and} \quad A'_{h+1} = \{\beta' \leq \theta \leq \pi\},$$

for suitable α, α', β and β' in $(0, \pi)$. It also implies $q_{m+1} = z_{s+1} - w_l = z'_{h+1} - w'_k$. For $t = 1, 2, 3, 4$, let $\theta_t \in [0, 2\pi)$ be such that $\{\theta = \theta_t\} = \mathcal{T}^{-1}\{\theta = (t-1)\pi/4\}$. The cones $A_s, -B_l, A'_h$ and $-B'_k$ are bounded by the rays $\{\theta = \theta_t\}$, by (1.2). By Remark 3.4, and since these cones are contained in $\{0 \leq \theta \leq \pi\}$, we may assume

$$(4.9) \quad 0 \leq \theta_1 < \theta_2 < \theta_3 < \theta_4 \leq \pi \quad \text{and} \quad \theta_4 \neq \theta_1 + \pi.$$

Moreover, the identities in (4.7) imply that one of the θ_t , (necessarily θ_1 , by (4.9)) equals 0. Assume $i = 1$ and $j = 2$ in (1.2). The condition (1.2), when expressed in terms of the θ_t , becomes

$$\{A_s, -B_l\} = \{\{0 \leq \theta \leq \theta_4\}, \{\theta_2 \leq \theta \leq \theta_3\}\} \quad \text{and} \\ \{A'_h, -B'_k\} = \{\{0 \leq \theta \leq \theta_2\}, \{\theta_3 \leq \theta \leq \theta_4\}\}.$$

Since (4.7) implies $A_s \neq \{\theta_2 \leq \theta \leq \theta_3\}$ and $A_h \neq \{\theta_3 \leq \theta \leq \theta_4\}$, we have

$$(4.10) \quad A_s = \{0 \leq \theta \leq \theta_4\}, \quad -B_l = \{\theta_2 \leq \theta \leq \theta_3\} \quad \text{and} \quad -B'_k = \{\theta_3 \leq \theta \leq \theta_4\}.$$

Similar arguments imply that if $i = 2$ and $j = 1$ in (1.2), then we have

$$(4.11) \quad A_s = \{0 \leq \theta \leq \theta_2\}, \quad -B_l = \{\theta_3 \leq \theta \leq \theta_4\} \quad \text{and} \quad -B'_k = \{\theta_2 \leq \theta \leq \theta_3\}.$$

Summarising, either (4.10) or (4.11) holds.

Let us prove that q_{m+1} is ambiguous, that is, since $q_{m+1} = z_{s+1} - w_l = z'_{h+1} - w'_k$, let us prove

$$(4.12) \quad \{A_{s+1}, -B_l\} \neq \{A'_{h+1}, -B'_k\}.$$

The cone $-B'_k$ does not belong to the set in the left-hand side of (4.12). Indeed, we have $\theta_3 < \theta_4 < \pi$, by (4.9) and the equality $\theta_1 = 0$. Thus (4.8), (4.10) and (4.11) imply $-B'_k \neq -B_l$ and $-B'_k \neq A_{s+1}$.

Since q_{m+1} is ambiguous, there exist a linear transformation $\mathcal{A}$ and $i, j \in \{1, 2\}$, with $i \neq j$, such that Alternative (ii) of Theorem 1.3 occurs, with $A = A_{s+1}$, $B = B_l$, $A' = A'_{h+1}$, $B' = B'_k$ and $\mathcal{T} = \mathcal{A}$. For $t = 1, 2, 3, 4$, let $\theta'_t \in [0, 2\pi)$ be such that $\{\theta = \theta'_t\} = \mathcal{A}^{-1}\{\theta = (t-1)\pi/4\}$. The θ'_t satisfy a condition analogous to (4.9) and, moreover, (4.8) implies $\theta'_4 = \pi$. It can be proved, by arguing as above, that one of the following possibilities occurs:

$$(4.13) \quad A_{s+1} = \{\theta'_1 \leq \theta \leq \pi\}, \quad -B_l = \{\theta'_2 \leq \theta \leq \theta'_3\} \quad \text{and} \quad -B'_k = \{\theta'_1 \leq \theta \leq \theta'_2\};$$

$$(4.14) \quad A_{s+1} = \{\theta'_3 \leq \theta \leq \pi\}, \quad -B_l = \{\theta'_1 \leq \theta \leq \theta'_2\} \quad \text{and} \quad -B'_k = \{\theta'_2 \leq \theta \leq \theta'_3\}.$$

Observe that (4.10) and (4.13) do not hold together, because (4.10) and (4.13) imply $\theta'_3 = \theta_3 = \theta'_1$, which contradicts $\theta'_3 > \theta'_1$. Similar arguments prove that (4.11) and (4.14) do not hold together. Assume (4.10) and (4.14). In this case we have $\theta_4 = \theta'_3$. Thus the identities in (4.11) and (4.14) regarding A_s and A_{s+1} become

$$A_s = \{0 \leq \theta \leq \theta_4\} \quad \text{and} \quad A_{s+1} = \{\theta_4 \leq \theta \leq \pi\}.$$

These conditions clearly imply that $[z_{s-1}, z_s]$ and $[z_{s+1}, z_{s+2}]$ are parallel to $\{\theta = \theta_4\}$. When (4.11) and (4.13) hold we have $\theta_2 = \theta'_1$. Thus we have $A_s = \{0 \leq \theta \leq \theta_2\}$ and $A_{s+1} = \{\theta_2 \leq \theta \leq \pi\}$, which again imply $[z_{s-1}, z_s]$ parallel to $[z_{s+1}, z_{s+2}]$. This concludes the proof if (4.5) holds and $[q_m, q_{m+1}] = [z'_h, z'_{h+1}] - w'_k$. If (4.5) holds and $[q_m, q_{m+1}] = z'_h - [w'_k, w'_{k+1}]$, then the claim can be proved as before, by substituting in the proof A'_h , A'_{h+1} and $-B'_k$ respectively with $-B'_k$, $-B'_{k+1}$ and A'_h . Similar arguments prove the claim if (4.5) is substituted by (4.6). $\square$

Claim 4.2.3. *The function $g_{K,L}$ determines $\{\text{cone}(K, K_u), \text{cone}(-L, (-L)_u)\}$ for each $u \in S^1$.*

Proof. Assume that no vertex of $K - L$ is ambiguous. Let $u \in S^1$. Claim 4.2.1 implies that $g_{K,L}$ distinguishes whether both K_u and L_{-u} are vertices, or both K_u and L_{-u} are edges or one is a vertex and the other one is an edge. If both K_u and L_{-u} are vertices, then the claim follows from the assumption that $(K - L)_u$ is not ambiguous. If K_u and L_{-u} are edges, then $\{\text{cone}(K, K_u), \text{cone}(-L, (-L)_u)\} = \{H\}$, where $H = \{x \in \mathbb{R}^2 : x \cdot u \leq 0\}$. If K_u is an edge and L_{-u} is a vertex (or vice versa), then consider the set

$$\bigcup_{w \in S^1: (K-L)_w \text{ is a vertex}} \{\text{cone}(K, K_w), \text{cone}(-L, (-L)_w)\}.$$

This set is determined by $g_{K,L}$, since no vertex of $K - L$ is ambiguous. The convexity of K and of L implies that only one among these cones, say A , has the property that $A \setminus \{O\} \subset \text{int } H$. Then

$$\{\text{cone}(K, K_u), \text{cone}(-L, (-L)_u)\} = \{H, A\}.$$

Assume that some vertex of $K - L$ is ambiguous. By Claim 4.2.2, all vertices are ambiguous. Let us use the notations introduced before Claim 4.2.2. Let s be any index in $\{1, \dots, n\}$ and choose $l \in \{1, \dots, t\}$ in such a way that $[z_s, z_{s+1}] - w_l$ is an edge of $K - L$. This is possible because if u denotes the outer normal to K at $[z_s, z_{s+1}]$, then L_{-u} is a vertex, by Claim 4.2.2. By the same claim, $[z_{s-1}, z_s]$ and $[z_{s+1}, z_{s+2}]$ are parallel. A similar argument proves that, given any $l \in \{1, \dots, t\}$, $[w_{l-1}, w_l]$ and $[w_{l+1}, w_{l+2}]$ are parallel. This is possible only if both K and L are parallelograms. Similar arguments prove that K' and L' are parallelograms too.

Consider a given vertex $q_m = z_s - w_l = z'_h - w'_k$ of $K - L$. Since q_m is ambiguous there exists a linear transformation $\mathcal{T}$ such that Alternative (ii) of Theorem 1.3 holds, with $A_s = A$, $B_l = B$, $A'_h = A'$ and $B'_k = B'$. Assume, for instance,

$$\mathcal{T}A_s = \mathcal{A}_1, \quad \mathcal{T}B_l = \mathcal{B}_1, \quad \mathcal{T}A'_h = \mathcal{A}_2 \quad \text{and} \quad \mathcal{T}B_l = \mathcal{B}_2.$$

Since $\mathcal{T}K$ is a parallelogram, the set of the directions of its edges coincides with the set of the directions of the edges of the support cone $\mathcal{T}A_s$ in one vertex of $\mathcal{T}K$. Thus the edges of $\mathcal{T}K$ are parallel to those of $\mathcal{K}_2$. Similarly, the edges of $\mathcal{T}L$ (of $\mathcal{T}K'$ and $\mathcal{T}L'$) are parallel to those of $\mathcal{L}_2$ (of $\mathcal{K}_1$ and $\mathcal{L}_1$, respectively). Let x_1 and x_2 be the centres of $\mathcal{T}K$ and of $\mathcal{T}K'$, respectively, and let y be the center of $\mathcal{T}L - x_1$ and $\mathcal{T}L' - x_2$ (these sets have equal center because $K - L = K' - L'$, by (2.2)). Choose the parameters defining $\mathcal{K}_2$ and $\mathcal{L}_2$ so that $\mathcal{T}K - x_1 = \mathcal{K}_2$ and $\mathcal{T}L - x_1 = \mathcal{L}_2 + y$. The edges of $\mathcal{T}K$ and $\mathcal{T}K'$ parallel to $\{\theta = 0\}$ have equal length, by Claim 4.2.1 and because $\mathcal{T}L$ and $\mathcal{T}L'$ have no edges parallel to $\{\theta = 0\}$. Also the edges of $\mathcal{T}K$ and $\mathcal{T}L'$ parallel to $\{\theta = 3\pi/4\}$ have equal length, and the same property holds for the edges of $\mathcal{T}K'$ and $\mathcal{T}L$ parallel to $\{\theta = \pi/4\}$, and for those of $\mathcal{T}L$ and $\mathcal{T}L'$ parallel to $\{\theta = \pi/2\}$. Therefore $\mathcal{T}K' - x_2 = \mathcal{K}_1$ and $\mathcal{T}L' - x_2 = \mathcal{L}_1 + y$. This contradicts the assumptions of Proposition 4.2 and proves the claim. Claim 4.2.1 and Claim 4.2.3 imply that $(K, -L)$ and $(K', -L')$ are synisothetic and conclude the proof of Proposition 4.2. $\square$

5. A CRUCIAL LEMMA

This section is dedicated to the proof of Lemma 5.5 and to some results needed in its proof. The first one, Lemma 5.1, is, in our opinion, of interest by itself. It is contained in the unpublished note [Man01], where it is proved with geometrical arguments. Here we present a different, shorter proof which is based on the Theorem of supports for convolutions [Hor83, Th. 4.3.3].

Lemma 5.1. *Let A , B , C and D be convex cones in $\mathbb{R}^n$, $n \geq 2$, with apex the origin O . Assume that each of them either coincides with $\{O\}$ or has non-empty interior and, moreover, $A \cup B \subset \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n : x_n \geq 0\}$, $A \cap \{x_n = 0\} = B \cap \{x_n = 0\} = \{O\}$, $C \cup D \subset \{x_n \leq 0\}$ and $\text{conv}(C \cup D)$ is pointed. If*

$$(5.1) \quad g_{A,C} + g_{B,D} = g_{A,D} + g_{B,C}$$

then either $A = B$ or $C = D$. The same conclusion holds if the hypothesis “ $\text{conv}(C \cup D)$ is pointed” is substituted by “either $A \subset B$ or $B \subset A$, and, moreover, either $C \subset D$ or $D \subset C$ ”.

Proof. For $r > 0$, let $A_r = A \cap B(O, r)$, $B_r = B \cap B(O, r)$, $C_r = C \cap B(O, r)$ and $D_r = D \cap B(O, r)$. We prove that there exists $s > 1$ such that, for each $x \in B(O, 1)$, we have

$$(5.2) \quad \begin{aligned} g_{A,C}(x) &= g_{A_s, C_s}(x), & g_{B,D}(x) &= g_{B_s, D_s}(x), \\ g_{B,C}(x) &= g_{B_s, C_s}(x) \text{ and } g_{A,D}(x) &= g_{A_s, D_s}(x). \end{aligned}$$

Since $A \cap \{x_n = 0\} = B \cap \{x_n = 0\} = \{O\}$, there exists $s > 1$ such that $(A \cup B) \cap \{x_n \leq 1\} \subset B(O, s-1)$. Let $x \in B(O, 1)$. Since $C + x \subset \{x_n \leq 1\}$, we have $A \cap (C + x) \subset A \cap \{x_n \leq 1\} \subset B(O, s-1)$. Moreover, by the triangle inequality, we have $(C + x) \cap B(O, s-1) \subset (C_s + x) \cap B(O, s-1)$. Therefore $A \cap (C + x) = A_s \cap (C_s + x)$ and $g_{A,C}(x) = g_{A_s, C_s}(x)$. Similar arguments prove the other identities in (5.2).

All the functions which appear in (5.1) are homogeneous of degree n and (5.1) holds true if and only if it holds true in $B(O, 1)$, that is, if and only if $g_{A_s, C_s}(x) + g_{B_s, D_s}(x) = g_{A_s, D_s}(x) + g_{B_s, C_s}(x)$ for each $x \in B(O, 1)$. By (1.1) this condition is equivalent to

$$(5.3) \quad (1_{A_s} - 1_{B_s}) * (1_{-C_s} - 1_{-D_s})(x) = 0 \quad \text{for each } x \in B(O, 1).$$

Let us conclude the proof under the assumption $\text{conv}(C \cup D)$ pointed. Let $S = \text{supp}(1_{A_s} - 1_{B_s}) * (1_{-C_s} - 1_{-D_s})$. By (5.3), we have $S \cap B(O, 1) = \emptyset$. The set S is clearly contained in $\text{conv}(A \cup B \cup (-C) \cup (-D))$ and the assumptions of the lemma imply that this union is pointed. Therefore the identity $S \cap B(O, 1) = \emptyset$ implies that there exists $\varepsilon > 0$ such that

$$(\text{conv } S) \cap B(O, \varepsilon) = \emptyset.$$

We may apply the Theorem of supports for convolutions [Hor83, Th. 4.3.3], since the involved functions have compact supports. This theorem implies

$$\text{conv } S = \text{conv supp}(1_{A_s} - 1_{B_s}) + \text{conv supp}(1_{-C_s} - 1_{-D_s}).$$

Therefore either we have $\text{conv supp}(1_{A_s} - 1_{B_s}) \cap B(O, \varepsilon/2) = \emptyset$ or we have $\text{conv supp}(1_{-C_s} - 1_{-D_s}) \cap B(O, \varepsilon/2) = \emptyset$. In the first case we have $A_s \cap B(O, \varepsilon/2) = B_s \cap B(O, \varepsilon/2)$, which is equivalent to $A = B$. In the second case, by similar arguments, we have $C = D$.

Drop the assumption $\text{conv}(C \cup D)$ pointed. When $A \subset B$ and $C \subset D$ then the functions which are convolved in (5.3) are constant in the interior of their supports, and their convolution vanish in $B(O, 1)$ if and only if one of them vanish. The other cases are treated similarly. $\square$




FIGURE 4. Up to affine transformations, $(\mathcal{K}_3, \mathcal{L}_3)$ and $(\mathcal{K}_4, \mathcal{L}_4)$ are the only pairs of convex polygons with equal cross covariogram which are synisothetic and are not trivial associates.

Let us introduce the second counterexample for the cross covariogram problem for convex polygons.

Example 5.2. Let α, β, γ and δ be positive real numbers, $m \in \mathbb{R}$, $y \in \mathbb{R}^2$, I_1 and I_3 be as in Example 4.1 and $I^{(m)} = (1/\sqrt{1+m^2})[(-m, -1), (m, 1)]$. Assume either $m = 0$, $\alpha \neq \gamma$ and $\beta \neq \delta$ or else $m \neq 0$ and $\alpha \neq \gamma$. We define four parallelograms as follows:

$$\mathcal{K}_3 = \alpha I_1 + \beta I_3; \quad \mathcal{L}_3 = \gamma I_1 + \delta I^{(m)} + y; \quad \mathcal{K}_4 = \gamma I_1 + \beta I_3; \quad \text{and} \quad \mathcal{L}_4 = \alpha I_1 + \delta I^{(m)} + y.$$

See Fig. 4. We have $g_{\mathcal{K}_3, \mathcal{L}_3} = g_{\mathcal{K}_4, \mathcal{L}_4}$ (it can be proved by arguing as in Example 4.1), and the pairs $(\mathcal{K}_3, -\mathcal{L}_3)$ and $(\mathcal{K}_4, -\mathcal{L}_4)$ are clearly synisothetic. However, $(\mathcal{K}_3, \mathcal{L}_3)$ and $(\mathcal{K}_4, \mathcal{L}_4)$ are not trivial associates.

Lemma 5.3. Let K, L, K' and L' be convex polygons satisfying the assumptions of Theorem 1.1 with $(K, -L)$ and $(K', -L')$ synisothetic. Assume that, for a given $u \in S^1$, K_u and K'_u are edges and have different lengths. Let I_K^1 (and I_K^2) be the edge of K adjacent to K_u that, in counterclockwise order on ∂K , precedes (and follows, respectively) K_u . For $i = 1, 2$, let I_{-L}^i , $I_{K'}^i$ and $I_{-L'}^i$ be respectively edges of $-L$, of K' and of $-L'$ defined in analogy to I_K^i . Then $(-L)_u$ and $(-L')_u$ are edges and

- (i) either I_K^i is parallel to I_{-L}^i and $I_{K'}^i$ is parallel to $I_{-L'}^i$, for $i = 1, 2$,
- (ii) or $I_K^1, I_K^2, I_{K'}^1$ and $I_{K'}^2$ are parallel and $I_{-L}^1, I_{-L}^2, I_{-L'}^1$ and $I_{-L'}^2$ are parallel.

Remark 5.4. When $(K, L) = (\mathcal{K}_3, \mathcal{L}_3)$, $(K', L') = (\mathcal{K}_4, \mathcal{L}_4)$ and $u = (0, 1)$, then Alternative (ii) of Lemma 5.3 occurs; see Fig. 4.

Proof. Since $\lambda_1(K_u) \neq \lambda_1(K'_u)$, the synisothesis of $(K, -L)$ and $(K', -L')$ implies (2.4). Therefore $(-L)_u$ and $(-L')_u$ are edges, $\lambda_1(K'_u) = \lambda_1((-L)_u)$ and $\lambda_1((-L')_u) = \lambda_1(K_u)$. Let $K_u = [z_0, z_1]$ and $(-L)_u = [w_0, w_1]$, where z_1 follows z_0 and w_1 follows w_0 , in counterclockwise order on the respective boundaries. For $i = 1, 2$, let $v_K^i \in S^1$ be parallel to I_K^i and oriented in such a way that $v_K^i \cdot u > 0$. Define $v_{-L}^i, v_{K'}^i$ and $v_{-L'}^i$ similarly. Assume, for instance, $\lambda_1(K_u) > \lambda_1(K'_u)$, that is, $\lambda_1(K_u) > \lambda_1((-L)_u)$ and $\lambda_1(K'_u) < \lambda_1((-L')_u)$. Let $q_0 = z_0 + w_0$, $q_1 = z_0 + w_1$, $q_2 = z_1 + w_0$ and $q_3 = z_1 + w_1$. The points q_0, q_1, q_2 and q_3 belong to $[q_0, q_3] = K_u + (-L)_u$. Moreover we have $[q_0, q_3] = (K - L)_u$ (by (2.1)) and $q_0 < q_1 < q_2 < q_3$ in counterclockwise order on $\partial(K - L)$.

Let $S^2(K, L) = \text{cl}\{x \in \mathbb{R}^2 : g_{K, L} \text{ is not } C^2 \text{ at } x\}$. We analyse the shape of $S^2(K, L) \cap W$, where W is a neighbourhood of $[q_0, q_3]$. It is easy to prove that

$$S^2(K, L) = \left(\bigcup_{z \text{ vertex of } K} (-\partial L + z) \right) \cup \left(\bigcup_{w \text{ vertex of } -L} (\partial K + w) \right).$$

Schmitt [Scm93] proves this formula when $K = L$ and the general case can be proved in the same way. We also recall Lemma 3.2, which proves the previous

formula when K and L are planar convex cones. Thus, when W is sufficiently small, we have

$$W \cap S^2(K, L) = W \cap \left((-\partial L + z_0) \cup (-\partial L + z_1) \cup (\partial K + w_0) \cup (\partial K + w_1) \right).$$

This set is the union of $[q_0, q_3]$ and, for each $i = 0, \dots, 3$, of two line segments (possibly coincident) containing q_i . If U_i denotes the set of the directions of the line segments containing q_i , then

$$U_0 = \{v_K^1, v_{-L}^1\}, U_1 = \{v_K^1, v_{-L}^2\}, U_2 = \{v_{K'}^2, v_{-L}^1\} \text{ and } U_3 = \{v_{K'}^2, v_{-L}^2\}.$$

The above analysis can be repeated for $g_{K', L'}$. However, in this case K' has the role of $-L$ and $-L'$ the role of K , because $\lambda_1(K'_u) < \lambda_1((-L)'_u)$. Therefore the identity $g_{K, L} = g_{K', L'}$ implies

$$U_0 = \{v_{K'}^1, v_{-L'}^1\}, U_1 = \{v_{K'}^2, v_{-L'}^1\}, U_2 = \{v_{K'}^1, v_{-L'}^2\} \text{ and } U_3 = \{v_{K'}^2, v_{-L'}^2\}.$$

Observe that if one of the equalities $v_K^1 = v_{-L'}^1$, $v_K^2 = v_{-L'}^2$, $v_{-L}^1 = v_{K'}^1$ and $v_{-L}^2 = v_{K'}^2$ holds, then also the other three equalities hold. For instance, if $v_K^1 = v_{-L'}^1$, then the identities involving U_0 imply $v_{-L}^1 = v_{K'}^1$, those involving U_1 imply $v_{-L}^2 = v_{K'}^2$. Once these are established, the identities involving U_2 imply $v_K^2 = v_{-L'}^2$. When one of these equalities holds, Alternative (i) occurs.

Assume $U_0 = U_1 = U_2 = \{v, w\}$, for suitable v, w , with $v \neq w$. If $v_K^1 = v_{-L'}^1$, then Alternative (i) occurs, as proved above. If $v_K^1 \neq v_{-L'}^1$ and, for instance, $v_K^1 = w$ and $v_{-L'}^1 = v$, then necessarily $v_K^1 = v_{K'}^2 = w$, $v_{-L}^1 = v_{-L}^2 = v$, $v_{K'}^1 = v_{K'}^2 = w$ and $v_{-L'}^1 = v_{-L'}^2 = v$, that is, Alternative (ii) occurs. When $U_0 = U_1 = U_2$ consists of a single element, clearly $v_K^1 = v_{-L'}^1$ and Alternative (i) occurs. If $U_0 \neq U_1$, then v_K^1 and $v_{-L'}^1$ coincide, because they are the only element of $U_0 \cap U_1$, and Alternative (i) occurs. Finally, if $U_0 \neq U_2$, then $v_{K'}^1$ and v_{-L}^1 coincide, because they are the only element of $U_0 \cap U_2$, and again Alternative (i) occurs. $\square$


Lemma 5.5. *Assume that K , K' , L and L' are convex polygons satisfying the assumptions of Theorem 1.1, that $(K, -L)$ and $(K', -L')$ are synisothetic and that (K, L) and (K', L') are not trivial associates. Assume also the following properties:*

- (i) *there exists an arc $U \subset S^1$, which is not a point, such that (2.3) holds true for each $u \in U$, and U is a maximal arc (with respect to inclusion) with this property;*
- (ii) *there exist $u_0 \in U$ such that $K_{u_0} = K'_{u_0}$ and $(-L)_{u_0} = (-L')_{u_0}$;*
- (iii) *if Σ denotes the maximal closed arc contained in $\partial K \cap \partial K'$ and containing K_{u_0} , and Ω denotes the maximal closed arc contained in $\partial(-L) \cap \partial(-L')$ and containing $(-L)_{u_0}$, then neither Σ nor Ω are points or line segments.*

Then Σ is a translate of Ω .

The proof of this lemma is divided in five steps and occupies all the rest of the section. First observe that $\Sigma \neq \partial K$ and $\Omega \neq \partial(-L)$. Indeed, if, for instance, $\Sigma = \partial K$, then $K = K'$. Moreover $L = L'$, because $K - L = K' - L'$ (by (2.2)) and the Minkowski addition satisfies a cancellation law (see [Sch93, p. 126]). Thus (K, L) and (K', L') are trivial associates, a contradiction.

Claim 5.5.1. *Let $a_1, a_2 \in \partial K$, $b_1, b_2 \in \partial(-L)$ and $u_1, u_2 \in S^1$ satisfy $\Sigma = [a_1, a_2]_{\partial K}$, $\Omega = [b_1, b_2]_{\partial(-L)}$ and $\text{cl } U = [u_1, u_2]_{S^1}$; see Fig. 5. Then, for each $j = 1, 2$, the sets K_{u_j} and K'_{u_j} are edges which contain a_j and have a common endpoint a'_j contained in $\text{relint } \Sigma$. Similarly, the sets $(-L)_{u_j}$ and $(-L')_{u_j}$ are edges which contain b_j and have a common endpoint b'_j contained in $\text{relint } \Omega$. In particular, Σ and $\Omega + a_j - b_j$ coincide in a neighbourhood of a_j .*

FIGURE 5. The sets U , Σ and Ω .

Moreover, if, for some $j \in \{1, 2\}$, either $\lambda_1(K_{u_j}) \neq \lambda_1(K'_{u_j})$ and Alternative (i) of Lemma 5.3 (with u replaced by u_j) holds, or else $\lambda_1(K_{u_j}) = \lambda_1(K'_{u_j})$, then

$$(5.4) \quad N(K, a_j) = N(-L', b_j) \quad \text{and} \quad N(K', a_j) = N(-L, b_j).$$

Proof. Let u_1^0 be the upper endpoint of $S^1 \cap N(K, a_1)$ and of $S^1 \cap N(K', a_1)$. These endpoints coincide because $[a_1, a_2]_{\partial K} \subset \partial K \cap \partial K'$. Define u_2^0 as u_1^0 , with lower replacing upper and a_2 replacing a_1 . Since $K_{u_0} = K'_{u_0} \subset \Sigma$ and K_{u_0} is isothetic to K'_{u_0} , by assumption, we have $u_0 \in [u_1^0, u_2^0]_{S^1}$. Let us prove

$$(5.5) \quad [u_1^0, u_2^0]_{S^1} = \text{cl } U.$$

If $u \in (u_1^0, u_2^0)_{S^1}$ then K_u and K'_u are isothetic, because K_u and K'_u are contained in $\text{relint } \Sigma$ and therefore ∂K and $\partial K'$ coincide in a neighbourhood of $K_u = K'_u$. The synisothesis of $(K, -L)$ and $(K', -L')$ implies that also $(-L)_u$ and $(-L')_u$ are isothetic. Thus (2.3) holds for each $u \in (u_1^0, u_2^0)_{S^1}$. Since $[u_1^0, u_2^0]_{S^1}$ intersects U (both arcs contains u_0) and U is maximal, we have $[u_1^0, u_2^0]_{S^1} \subset \text{cl } U$.

In order to conclude the proof of (5.5) it suffices to show that in any neighbourhood of u_j^0 there are directions u for which (2.3) does not hold, for $j = 1, 2$. Since ∂K and $\partial K'$ bifurcate at a_j , a_j is a vertex of K or of K' . If it is both a vertex of K and of K' then the vertex a_j of K is not isothetic to the vertex a_j of K' . In this case (2.3) does not hold for all $u \in N(K, a_j) \cup N(K', a_j)$. If a_j is a vertex of one polygon and it belongs to the relative interior of an edge of the other polygon, then the unit outer normal to this edge is necessarily u_j^0 . Since ∂K and $\partial K'$ bifurcate at a_j , (2.3) does not hold when $u = u_j^0$.

The identity (5.5) implies $u_1 = u_1^0$ and $u_2 = u_2^0$. The definition of u_j^0 clearly implies that K_{u_j} and K'_{u_j} are line segments, for $j = 1, 2$, and that $K_{u_j} \cap K'_{u_j}$ is a line segment which, in counterclockwise order on ∂K , follows a_j when $j = 1$ and precedes a_j when $j = 2$. Assumption (iii) of Lemma 5.5 implies that the upper endpoints of K_{u_1} and K'_{u_1} coincide and that this point belongs to $\text{relint } \Sigma$. Similar arguments prove the analogous property for the lower endpoints of K_{u_2} and K'_{u_2} . The properties proved up till now for K and K' can be proved, by similar arguments, also for $-L$ and $-L'$.

If, for some j , $\lambda_1(K_{u_j}) \neq \lambda_1(K'_{u_j})$, then (5.4) is an immediate consequence of Alternative (i) of Lemma 5.3. Assume $\lambda_1(K_{u_j}) = \lambda_1(K'_{u_j})$. In this case the synisothesis of $(K, -L)$ and $(K', -L')$ implies $\lambda_1((-L)_{u_j}) = \lambda_1((-L')_{u_j})$. Therefore, a_j is a vertex of K and of K' and b_j is a vertex of $-L$ and of $-L'$. Since $N(K, a_j) \cap S^1$, $N(K', a_j) \cap S^1$, $N(-L, b_j) \cap S^1$ and $N(-L', b_j) \cap S^1$ have the endpoint u_j in common, there exists a ‘‘perturbation’’ $\bar{u}$ of u_j which belongs to the relative interior of $N(K, a_j)$, $N(K', a_j)$, $N(-L, b_j)$ and $N(-L', b_j)$. The vertex a_j of K is not isothetic to the vertex a_j of K' , because ∂K and $\partial K'$ bifurcate at a_j . Therefore, the vertex a_j of K is isothetic to the vertex b_j of $-L'$, and the vertex a_j of K' is isothetic to the vertex b_j of $-L$. This is equivalent to (5.4). $\square$




FIGURE 6. $K \cap (L + x_0)$ (dotted lines) and $K' \cap (L' + x_0)$ (continuous lines).

Let $\mathcal{R}$ denote clockwise rotation by $\pi/2$ and let

$$(5.6) \quad \theta = \mathcal{R} \left(\frac{a_2 - a_1}{\|a_2 - a_1\|} \right) \quad \text{and} \quad \theta' = \mathcal{R} \left(\frac{b_2 - b_1}{\|b_2 - b_1\|} \right).$$

Since the assumptions and the conclusion of Lemma 5.5 are preserved by substituting K , $-L$, K' and $-L'$ with $-L$, K , $-L'$ and K' , respectively, we may assume $\theta \leq \theta'$ without loss of generality. The next claim states that, under suitable hypotheses, the arcs Σ and $\Omega + a_1 - b_1$ can bifurcate only at a point c_1 where every outer normal to these arcs is θ , or it is larger than θ' .

Claim 5.5.2. *Let θ and θ' be as in (5.6), with $\theta \leq \theta'$. For $i = 1, 2$, let a_i, a'_i, b_i, b'_i and u_i be as in Claim 5.5.1 and let $x_i = b_i - a_i$. Assume that for some $j \in \{1, 2\}$ the following two hypotheses hold: if $\lambda_1(K_{u_j}) \neq \lambda_1(K'_{u_j})$, then Alternative (i) of Lemma 5.3, with u replaced by u_j , holds; we have $c_j \in \text{relint } \Sigma \cap \text{relint } (\Omega + x_j)$, where c_j is the endpoint different from a_j of the arc that is contained in $\Sigma \cap (\Omega + x_j)$ and contains a_j . Then, when $j = 1$ we have*

$$(5.7) \quad N(K, c_1) \cap N(-L + x_1, c_1) \cap [u_1, \theta]_{S^1} = \emptyset \quad \text{and}$$

$$(5.8) \quad N(K, c_1) \cap N(-L + x_1, c_1) \cap (\theta, \theta')_{S^1} = \emptyset,$$

while when $j = 2$ we have

$$(5.9) \quad N(K, c_2) \cap N(-L + x_2, c_2) \cap (\theta', u_2]_{S^1} = \emptyset \quad \text{and}$$

$$(5.10) \quad N(K, c_2) \cap N(-L + x_2, c_2) \cap (\theta, \theta')_{S^1} = \emptyset.$$

Proof. We prove the claim when $j = 1$. To prove (5.7) assume that there exists $w \in S^1$ in the intersection of the three sets. Let us first prove that a_1 is a vertex of K and of K' when $w = u_1$. Indeed, due to Claim 5.5.1, a_1 is a vertex of both sets if and only if $\lambda_1(K_{u_1}) = \lambda_1(K'_{u_1})$. If $\lambda_1(K_{u_1}) \neq \lambda_1(K'_{u_1})$, then Alternative (i) of Lemma 5.3 implies that K and $-L' + x_1$ coincide in a neighbourhood of K_{u_1} . In particular, $c_1 \notin [a_1, a'_1]$. Thus $u_1 \notin N(K, c_1)$, a contradiction to $w = u_1$. Similar arguments prove that b_1 is a vertex of $-L$ and of $-L'$ when $w = u_1$.

Let $x_0 = a_1 + c_1 - x_1$. We claim that $g_{K,L}(x) \neq g_{K',L'}(x)$ for some x close to x_0 . First we prove

$$(5.11) \quad K \cap (L + x_0) = K' \cap (L' + x_0).$$

The translation by x_0 maps the points $-c_1 + x_1$ and $-b_1$ of $\partial L \cap \partial L'$ respectively to a_1 and c_1 ; see Fig. 6. Let $\pi = \{y \in \mathbb{R}^2 : (y - a_1) \cdot w \geq 0\}$. The sets $K, K', -L + x_1$ and $-L' + x_1$ are contained in $-\pi + x_1 + x_0 = \{y \in \mathbb{R}^2 : (y - c_1) \cdot w \leq 0\}$ (because w is an outer normal to all these sets at c_1). The inclusions $-L + x_1, -L' + x_1 \subset -\pi + x_1 + x_0$ are equivalent to $L + x_0, L' + x_0 \subset \pi$. Therefore

$$(5.12) \quad K \cap (L + x_0), K' \cap (L' + x_0) \subset \pi \cap (-\pi + x_1 + x_0).$$




FIGURE 7. $K \cap (L+x)$ (dotted lines) and $K' \cap (L'+x)$ (continuous lines) for a suitable x close to x_0 .

We claim that

$$(5.13) \quad K \cap \pi = K' \cap \pi \quad \text{and} \quad \text{cl}(K \Delta K') \cap \pi = \{a_1\}.$$

When $w = u_1$ the first identity is true because $K \cap \pi = [a_1, a'_1] = K' \cap \pi$, by Claim 5.5.1 and because a_1 is a vertex of K and of K' . When $w \in (u_1, \theta)_{S^1}$, we have $\partial K \cap \pi = \partial K' \cap \pi$, because $\partial K \cap \pi$ and $\partial K' \cap \pi$ are contained in Σ (observe that $a_2 \notin \pi$ because $w < \theta$), which is contained in $\partial K \cap \partial K'$. Since K and K' are convex, the identity $\partial K \cap \pi = \partial K' \cap \pi$ implies $K \cap \pi = K' \cap \pi$. To prove the second identity observe that $K \Delta K'$ is contained in one of the halfplanes bounded by the line through a_1 and a_2 , and observe that this halfplane intersects $\pi \cap K$ only in a_1 . Similar arguments prove $(-L + x_1) \cap \pi = (-L' + x_1) \cap \pi$ and $\text{cl}((-L + x_1) \Delta (-L' + x_1)) \cap \pi = \{a_1\}$. These identities are equivalent to

$$(5.14) \quad (L + x_0) \cap (-\pi + x_1 + x_0) = (L' + x_0) \cap (-\pi + x_1 + x_0) \quad \text{and}$$

$$(5.15) \quad \text{cl}((L + x_0) \Delta (L' + x_0)) \cap (-\pi + x_1 + x_0) = \{c_1\}.$$

Formulas (5.12), (5.13) and (5.14) imply (5.11).

Formulas (5.13) and (5.15) imply that $\text{cl}((K \Delta K') \cup ((L + x_0) \Delta (L' + x_0)))$ intersects the strip $\pi \cap (-\pi + x_1 + x_0)$ only in a_1 and c_1 . Therefore, when x is close to x_0 , the sets $K \cap (L+x)$ and $K' \cap (L'+x)$ may differ only in a neighbourhood of a_1 and of c_1 (see Fig. 7). Assume $K \subset K'$ in a neighbourhood of a_1 (note that either we have $K \subset K'$ or we have $K' \subset K$, because ∂K and $\partial K'$ coincide on one side of a_1). Define $A = \text{cone}(K', a_1) \setminus \text{cone}(K, a_1)$. By (5.4) we have $A = \text{cone}(-L, b_1) \setminus \text{cone}(-L', b_1)$. Let $C = \text{cone}(L, x_1 - c_1) = \text{cone}(L', x_1 - c_1)$ and $-D = \text{cone}(K, c_1) = \text{cone}(K', c_1)$ (recall that $K = K'$ near c_1 and $L = L'$ near $x_1 - c_1$). If $\delta > 0$ is chosen sufficiently small, outside $B(a_1, \delta) \cup B(c_1, \delta)$ we have $K \cap (L+x) = K' \cap (L'+x)$. On the other hand

$$((K' \cap (L' + x)) \setminus (K \cap (L + x))) \cap B(a_1, \delta) = (A + a_1) \cap (C + x_1 - c_1 + x);$$

see Fig. 7. Therefore

$$\begin{aligned} \lambda_2(K \cap (L+x) \cap B(a_1, \delta)) - \lambda_2(K' \cap (L'+x) \cap B(a_1, \delta)) \\ &= -\lambda_2((A + a_1) \cap (C + x_1 - c_1 + x)) \\ &= -\lambda_2(A \cap (C + x - x_0)) \\ &= -g_{A,C}(x - x_0). \end{aligned}$$

Similar arguments prove

$$\lambda_2(K \cap (L+x) \cap B(c_1, \delta)) - \lambda_2(K' \cap (L'+x) \cap B(c_1, \delta)) = g_{A,D}(x - x_0).$$

Therefore, for each x in a neighbourhood of x_0 , we have

$$(5.16) \quad g_{K,L}(x) - g_{K',L'}(x) = g_{A,D}(x - x_0) - g_{A,C}(x - x_0).$$

We apply Lemma 5.1, with $B = \{O\}$, $n = 2$ and the Cartesian coordinates chosen so that $w = (0, -1)$. The assumptions of this lemma are satisfied. Indeed, $C, D \subset \{x_2 \leq 0\}$, and either $C \subset D$ or $D \subset C$, because the lower endpoints of $C \cap S^1$ and of $D \cap S^1$ coincide, by $[a_1, c_1]_{\partial K} \subset \partial K \cap \partial(-L + x_1)$. Moreover, $\text{cl}(K \Delta K') \cap \pi = \{a_1\}$ implies $A \cap \{x_2 = 0\} = \{O\}$. Observe that we have $C \neq D$, because ∂K and $\partial(-L) + x_1$ bifurcate at c_1 . Thus, this lemma implies $g_{A,D} \neq g_{A,C}$. Since $g_{A,D}$ and $g_{A,C}$ are homogeneous functions of degree 2, they do not coincide in any neighbourhood of O . Thus (5.16) implies $g_{K,L} \neq g_{K',L'}$. This contradiction proves (5.7).

The proof of (5.8) is similar although simpler. Assume that $w \in S^1$ belongs to the three sets in (5.8) and define $x_0 = a_2 + c_1 - x_1$. We prove again that $g_{K,L}$ does not coincide with $g_{K',L'}$ in a neighbourhood of x_0 . The translation by x_0 maps the point $-c_1 + x_1$ of $\partial L \cap \partial L'$ to the point a_2 . The identity $K \cap (L + x_0) = K' \cap (L' + x_0)$ is proved as before. Defining $\pi = \{y : (y - a_2) \cdot w \geq 0\}$, the inclusion (5.12) holds also in this case. Moreover, when x is close to x_0 the sets $K \cap (L + x)$ and $K' \cap (L' + x)$ may differ only in a neighbourhood of a_2 , because $\text{cl}(K \Delta K')$ intersects the strip $\pi \cap (-\pi + x_1 + x_0)$ only in a_2 (recall that $a_1 \notin \pi$ because $w > \theta$) and $\text{cl}((L + x_0) \Delta (L' + x_0))$ does not intersect this strip (recall that $-b_1 + x_0 \notin -\pi + x_1 + x_0$ because this is equivalent to $a_1 \notin \pi$ which holds true, and $-b_2 + x_0 \notin -\pi + x_1 + x_0$ because $w < \theta'$). Assume $K \subset K'$ in a neighbourhood of a_2 , let $A = \text{cone}(K', a_2) \setminus \text{cone}(K, a_2)$ and let $C = \text{cone}(L, x_1 - c_1) = \text{cone}(L', x_1 - c_1)$. For each x close to x_0 we have

$$g_{K,L}(x) - g_{K',L'}(x) = -g_{A,C}(x - x_0).$$

Lemma 5.1, with $D = B = \{O\}$, implies that the previous formula contradicts $g_{K,L} = g_{K',L'}$. This contradiction proves (5.8). $\square$

Claim 5.5.3. *Let $\text{cl}U = [u_1, u_2]$. If $\lambda_1(K_{u_j}) \neq \lambda_1(K'_{u_j})$, for some $j \in \{1, 2\}$, then Alternative (i) of Lemma 5.3, with u substituted by u_j , holds.*

Proof. Let θ and θ' be as in (5.6). We may assume $\theta \leq \theta'$. For $i = 1, 2$, let a_i, a'_i, b_i and b'_i be as in Claim 5.5.1 and let $x_i = b_i - a_i$. Assume the claim false for some j . By (2.4), with u substituted by u_j , we have


$$(5.17) \quad \lambda_1(K_{u_j}) = \lambda_1((-L')_{u_j}) \quad \text{and} \quad \lambda_1(K'_{u_j}) = \lambda_1((-L)_{u_j}).$$

Let a''_j be the endpoint of K_{u_j} which does not belong to K'_{u_j} or the endpoint of K'_{u_j} which does not belong to K_{u_j} , according to whether $\lambda_1(K_{u_j}) > \lambda_1(K'_{u_j})$ or $\lambda_1(K_{u_j}) < \lambda_1(K'_{u_j})$. Define b''_j similarly, by substituting K with $-L$ and K' with $-L'$.

First we prove that if Claim 5.5.4 is false both when $j = 1$ and when $j = 2$, then $(K, -L)$ and $(K', -L')$ are not synisothetic, contrary to what assumed in Lemma 5.5. Let $w \in S^1$ be orthogonal to the edges of K adjacent to K_{u_2} and to those of K' adjacent to K'_{u_2} , oriented in such a way that $w \in N(K, a'_2)$. We recall that Lemma 5.3 proves that those edges are parallel. Define v as w , with K, K' and a'_2 replaced by $-L, -L'$ and b'_2 , respectively. We stress that $w \neq v$, because otherwise both (i) and (ii) in Lemma 5.3 hold true, and this implies that Claim 5.5.4 is true when $j = 2$. We distinguish two possible cases.

Case $a'_1 = a'_2$. We have $\Sigma = [a_1, a'_1] \cup [a'_2, a_2]$, $w = u_1$ and $u_2 \in (u_1, -u_1)_{S^1}$; see Fig. 8.

Case $a'_1 \neq a'_2$. We have $\Sigma = [a_1, a'_1] \cup [a'_1, a'_2] \cup [a'_2, a_2]$, $w \neq u_1$ and $w \neq u_2$. Moreover, the parallelism of the edges of K adjacent to K_{u_i} and the parallelism of

FIGURE 8. The two possible configurations for K and K' .

the edges of K' adjacent to K'_{u_i} , for each $i \in \{1, 2\}$, together with the convexity of K and K' , imply that $K = \text{conv}(a_1, a'_1, a'_2, a_2)$ and $K' = \text{conv}(a''_1, a'_1, a'_2, a''_2)$, or vice versa.

Similar descriptions hold also for $-L$ and $-L'$, with v , b_i , b'_i and b''_i replacing, respectively, w , a_i , a'_i and a''_i .

It cannot be $a'_1 = a'_2$ and $b'_1 = b'_2$, because the descriptions above imply $w = v$, which is false. Assume $a'_1 \neq a'_2$ and $b'_1 \neq b'_2$. If, say, $K' = \text{conv}(a''_1, a'_1, a'_2, a''_2)$, then no edge of $-L$ is a translate of the edge $[a''_1, a''_2]$ of K' , because no edge of $-L$ is orthogonal to w . Thus, the synisothesis of $(K, -L)$ and $(K', -L')$ implies that an edge of K is a translate of $[a''_1, a''_2]$. This implies that $u_2 = -u_1$. Both K , K' , L and L' are parallelograms with two edges orthogonal to u_1 . This property and (5.17) imply that, up to an affine transformation, (K, L) and (K', L') are trivial associates of $(\mathcal{K}_3, \mathcal{L}_3)$ and $(\mathcal{K}_4, \mathcal{L}_4)$, respectively. This contradicts the assumptions of Lemma 5.5.

Assume $a'_1 \neq a'_2$ and $b'_1 = b'_2$. We have $\lambda_1([a''_1, a''_2]) > \lambda_1([a_1, a_2]) > \lambda_1([a'_1, a'_2])$, because $u_2 \in (u_1, -u_1)_{S^1}$. We also have $\lambda_1([a''_1, a''_2]) = \lambda_1([b''_1, b''_2]) + \lambda_1([a'_1, a'_2]) > \lambda_1([b'_1, b'_2]) > \lambda_1([b_1, b_2])$, by the previous descriptions, (5.17) with $j = 1$ and (5.17) with $j = 2$. Therefore, if, say, $K' = \text{conv}(a''_1, a'_1, a'_2, a''_2)$, then neither K nor $-L$ have an edge which is a translate of the edge $[a''_1, a''_2]$ of K' . This contradicts the synisothesis of $(K, -L)$ and $(K', -L')$.

It remains to prove that it is not possible that Claim 5.5.4 holds for one index, say $j = 1$, and it does not hold for the other one, say $j = 2$. Assume this false and let w and v be defined as above. Up to exchanging the roles of K and K' we may assume $\lambda_1(K_{u_2}) < \lambda_1(K'_{u_2})$, so that a_2 is a vertex of K and $N(K, a_2) \cap S^1 = [u_2, -w]_{S^1}$. Therefore, since $[a_1, a_2]$ is a chord of K , we have

$$(5.18) \quad w \leq \theta \text{ and } w = \theta \text{ if and only if } [a_1, a_2] \text{ is an edge of } K.$$

Similar arguments prove that

$$(5.19) \quad v \leq \theta' \text{ and } v = \theta' \text{ if and only if } [b_1, b_2] \text{ is an edge of } -L'.$$

We claim that $a'_2 \notin [a_1, c_1]_{\partial K}$. If $a'_2 \in [a_1, c_1]_{\partial K}$, then ∂K and $\partial(-L') + x_1$ coincide in a neighbourhood of a'_2 . In particular $-L'$ has a vertex p with $N(-L', p) \cap S^1 = [w, u_2]_{S^1}$. This is false, because b'_2 is the only vertex of $-L'$ with the property that u_2 is the upper endpoint of the intersection of S^1 with the normal cone at that vertex, however the lower endpoint of $N(-L, b'_2)$ is v and $w \neq v$. This contradiction proves $a'_2 \notin [a_1, c_1]_{\partial K}$.

Let z be the lower endpoint of $N(K, c_1) \cap S^1$. The vector z is also the lower endpoint of $N(-L' + x_1, c_1) \cap S^1$, since $[a_1, c_1]_{\partial K} \subset \partial K \cap (\partial(-L') + x_1)$. Since $a'_2 \notin [a_1, c_1]_{\partial K}$ we have $z \leq w$, by convexity. Thus, (5.7) and (5.18) imply $z = w = \theta$. The line segment $[a_1, a_2]$ is an edge of K , by (5.18). Therefore, by (5.4) with $j = 1$, the line l through b_1 and orthogonal to θ supports $-L'$. This implies $\theta \geq \theta'$. Since we assumed $\theta \leq \theta'$, we have $\theta = \theta'$. Thus $b_2 \in l$ and $[b_1, b_2]$ is an edge of $-L'$. The equalities $\theta' = v$ (a consequence of (5.19)), $\theta = w$ and $\theta = \theta'$ contradict $v \neq w$. $\square$

Claims 5.5.1 and 5.5.3 imply that (5.4) holds both when $j = 1$ and when $j = 2$.

Claim 5.5.4. *Let $\text{cl } U = [u_1, u_2]$ and, for $i = 1, 2$, let $x_i = a_i - b_i$, where a_i and b_i are as in Claim 5.5.1. Assume that Σ is not a translate of Ω . Then U contains an half-circle and $x_1 \neq x_2$. Moreover, if U is an half-circle, then Σ is the union of three consecutive edges L_1 , M and L_2 of a parallelogram, L_1 and L_2 are orthogonal to u_1 and Ω is the union of $L_1 - x_1$, of a line segment M' parallel to M and of $L_2 - x_2$.*

Proof. Let θ and θ' be as in (5.6) and, for $i = 1, 2$, let a'_i and b'_i be as in Claim 5.5.1 and let c_i be as in Claim 5.5.2. We may assume $\theta \leq \theta'$. First we prove

$$(5.20) \quad c_i \in \text{relint } \Sigma \cap (\text{relint } \Omega + x_i),$$

for $i = 1, 2$. Assume (5.20) false when $i = 2$. Since Σ and $\Omega + x_2$ coincide in a neighbourhood of their common upper endpoint a_2 , by Claim 5.5.1, c_2 coincides with the lower endpoint of one arc, that is, with a_1 or with $b_1 + x_2$. Assume $c_2 = a_1$. In this case $\Sigma \subset \Omega + x_2$ and, since $\Sigma \neq \Omega + x_2$ by assumption, $a_1 \neq b_1 + x_2$. The portion of Ω with outer normal u_1 , that is $[a_1, a'_1]$, should be contained in the portion of $\Omega + x_2$ with outer normal u_1 , that is in $[b_1, b'_1] + x_2$. Moreover, the inequality $a_1 \neq b_1 + x_2$ implies $\lambda_1([a_1, a'_1]) < \lambda_1([b_1, b'_1])$. The previous description implies that $\Omega + x_1$ bifurcates from Σ at a'_1 , that is, it implies $c_1 = a'_1$. Claim 5.5.3 and the observation that $c_1 = a'_1 \in \text{relint } \Sigma \cap (\text{relint } \Omega + x_1)$ imply that (5.7) holds. On the other hand, $c_1 = a'_1$ implies $u_1 \in N(K, c_1) \cap N(-L + x_1, c_1)$, which contradicts (5.7). Similar arguments prove $c_2 \neq b_1 + x_2$, and prove (5.20) when $i = 1$.

Let w_1 be the lower endpoint of $N(K, c_1) \cap S^1$ and of $N(-L' + x_1, c_1) \cap S^1$ (these endpoints coincide because $[a_1, c_1]_{\partial K} \subset \Omega \cap (\partial(-L) + x_1)$). Let w_2 be the upper endpoint of $N(K, c_2) \cap S^1$ and $N(-L' + x_2, c_2) \cap S^1$. We prove

$$(5.21) \quad w_1 \leq w_2.$$

Assume $w_1 > w_2$. In this case the two sub-arcs $[a_1, c_1]_{\partial K}$ and $[c_2, a_2]_{\partial K}$ of ∂K overlap and contain the arc $[c_2, c_1]_{\partial K}$. Moreover $[c_2, c_1]_{\partial K}$ is neither a point nor a line segment, because it contains a segment orthogonal to w_i , for $i = 1, 2$, by definition of w_i . Therefore the inclusions

$$[c_2, c_1]_{\partial K} \subset [a_1, c_1]_{\partial K} \subset \Omega + x_1 \quad \text{and} \quad [c_2, c_1]_{\partial K} \subset [c_2, a_2]_{\partial K} \subset \Omega + x_2,$$

imply $x_1 = x_2$. Since $\Omega + x_1$ is contained in the union of two overlapping sub-arcs of Σ , and its endpoints coincide with those of Σ , we have $\Omega + x_1 = \Sigma$. This contradicts the assumptions of Claim 5.5.4 and proves (5.21).

Claim 5.5.3 and (5.20) imply that the assumptions of Claim 5.5.2 are satisfied for each $j = 1, 2$. Thus (5.7), (5.8), (5.9) and (5.10) hold.

Assume $\theta = \theta'$. Formulas (5.7) and (5.9) imply respectively $w_1 \geq \theta$ and $w_2 \leq \theta$. Thus (5.21) implies $w_1 = \theta = w_2$. Let $K_\theta = [\dot{a}_1, \dot{a}_2]$ and $(-L)_\theta = [\dot{b}_1, \dot{b}_2]$, where $\dot{a}_2$ follows $\dot{a}_1$ and $\dot{b}_2$ follows $\dot{b}_1$, in counterclockwise order on the respective boundaries. We may clearly write

$$(5.22) \quad \begin{aligned} \Sigma &= [a_1, \dot{a}_1]_{\partial K} \cup [\dot{a}_1, \dot{a}_2] \cup [\dot{a}_2, a_2]_{\partial K} \quad \text{and} \\ \Omega &= [b_1, \dot{b}_1]_{\partial(-L)} \cup [\dot{b}_1, \dot{b}_2] \cup [\dot{b}_2, b_2]_{\partial(-L)}, \end{aligned}$$

where $[\dot{a}_1, \dot{a}_2]$ and $[\dot{b}_1, \dot{b}_2]$ are parallel. The definitions of w_1 and w_2 imply $c_i \in [\dot{a}_1, \dot{a}_2]$ and $c_i \in [\dot{b}_1, \dot{b}_2] + x_i$, for $i = 1, 2$, $c_1 \neq \dot{a}_1$ and $c_2 \neq \dot{a}_2$. Therefore, by definition of c_1 and of c_2 we have

$$(5.23) \quad [b_1, \dot{b}_1]_{\partial(-L)} = [a_1, \dot{a}_1]_{\partial K} - x_1 \quad \text{and} \quad [\dot{b}_2, b_2]_{\partial(-L)} = [\dot{a}_2, a_2]_{\partial K} - x_2.$$

We have $\lambda_1([\dot{a}_1, \dot{a}_2]) \neq \lambda_1([\dot{b}_1, \dot{b}_2])$, because otherwise $\Sigma = \Omega + x_1$, by (5.22) and (5.23).




FIGURE 9. $K \cap (L+x)$ (dotted lines) and $K' \cap (L'+x)$ (continuous lines) when $x = x_0$ (left) and when $x = x_0 - \varepsilon\theta$ (right).

In order to prove that U contains an half-circle it suffices to prove

$$(5.24) \quad \|a_2 - a_1\| \leq \|\dot{a}_2 - \dot{a}_1\|.$$

Indeed, if U does not contain an half-circle, then the line l through a_1 orthogonal to u_1 and the line r through a_2 orthogonal to u_2 bound a cone with apex contained in the halfplane bounded by the line through a_1 and a_2 and containing $[\dot{a}_1, \dot{a}_2]$. Since l and r support K , the cone contains $[\dot{a}_1, \dot{a}_2]$ and (5.24) is false.

We assume $\|a_2 - a_1\| > \|\dot{a}_2 - \dot{a}_1\|$ and obtain a contradiction by proving that $g_{K,L}(x) \neq g_{K',L'}(x)$, for some x close to x_0 , where x_0 will be defined later. We may write

$$a_2 - a_1 = \alpha(\dot{a}_2 - \dot{a}_1) \quad \text{and} \quad \dot{b}_2 - \dot{b}_1 = \beta(\dot{a}_2 - \dot{a}_1),$$

with $\alpha > 1$, $\beta > 0$. The inequality $\lambda_1([\dot{a}_1, \dot{a}_2]) \neq \lambda_1([\dot{b}_1, \dot{b}_2])$ implies $\beta \neq 1$. Up to exchanging K, K', a_i and $\dot{a}_i$ with $-L, -L', b_i$ and $\dot{b}_i$, respectively, we may assume $\beta < 1$. Let $x_0 = \dot{a}_1 + b_1 + \gamma(\dot{a}_2 - \dot{a}_1)$, where γ is a fixed number in $(\beta, \min(1, \beta + \alpha - 1))$. Easy computations and the equalities $a_2 - \dot{a}_2 = b_2 - \dot{b}_2$ and $a_1 - \dot{a}_1 = b_1 - \dot{b}_1$ (that are consequences of (5.23)) give the following expressions: $-b_1 + x_0 = \dot{a}_1 + \gamma(\dot{a}_2 - \dot{a}_1)$; $-\dot{b}_1 + x_0 = a_1 + (\gamma/\alpha)(a_2 - a_1)$; $-\dot{b}_2 + x_0 = a_1 + ((\gamma - \beta)/\alpha)(a_2 - a_1)$ and $-b_2 + x_0 = \dot{a}_1 + (1 - \alpha + \gamma - \beta)(\dot{a}_2 - \dot{a}_1)$. Since $0 < (\gamma - \beta)/\alpha < \gamma/\alpha < 1$ and $1 - \alpha + \gamma - \beta < 0$, the previous expressions imply the following formulas:

$$(5.25) \quad -\dot{b}_1 + x_0, -\dot{b}_2 + x_0 \in \text{relint}[a_1, a_2];$$

$$(5.26) \quad -b_2 + x_0 \notin K \cup K';$$

$$(5.27) \quad -b_1 + x_0 \in \text{relint}[\dot{a}_1, \dot{a}_2];$$

$$(5.28) \quad \|-b_1 + x_0 - \dot{a}_1\| > \|\dot{b}_2 - \dot{b}_1\|.$$

See Fig. 9. Let $\pi = \{y \in \mathbb{R}^2 : (y - a_1) \cdot \theta \geq 0\}$. The halfplane π contains Σ and its boundary contains a_1, a_2 and $b_1 + x_1$. Since $\theta = \theta'$, we have $b_2 + x_1 \in \partial\pi$ and $\Omega + x_1 \subset \pi$. By convexity of the involved polygons, $K \Delta K'$ and $(-L + x_1) \Delta (-L' + x_1)$ are contained in $\mathbb{R}^2 \setminus \pi$. The halfplane $-\pi + x_1 + x_0$ has outer normal θ and its boundary contains $\dot{a}_1, \dot{a}_2$ and c_1 . Thus $-\pi + x_1 + x_0$ contains $K, K', -L + x_1$ and $-L' + x_1$, since its boundary support the four sets at c_1 . Summarising, the following inclusions hold for any $x \in \mathbb{R}^2$:

$$(5.29) \quad \begin{aligned} K, K' &\subset -\pi + x_1 + x_0; & L + x, L' + x &\subset \pi - x_0 + x; \\ K \Delta K' &\subset \mathbb{R}^2 \setminus \pi; & (L + x) \Delta (L' + x) &\subset \mathbb{R}^2 \setminus (-\pi + x_1 + x). \end{aligned}$$

Therefore $K \cap (L + x)$ and $K' \cap (L' + x)$ are contained in the strip $N_1(x) = (\pi - x_0 + x) \cap (-\pi + x_1 + x_0)$, while $K \Delta K'$ and $(L + x) \Delta (L' + x)$ do not intersect

the strip $N_2(x) = \pi \cap (-\pi + x_1 + x)$. Since $N_1(x_0) = N_2(x_0)$ we have $K \cap (L + x_0) = K' \cap (L' + x_0)$. Let $x = x_0 - \varepsilon\theta$, with $\varepsilon > 0$. We have

$$(5.30) \quad (K \cap (L + x)) \Delta (K' \cap (L' + x)) \subset N_1(x) \setminus N_2(x) = \dot{\pi} \cup \ddot{\pi},$$

where $\dot{\pi} = (\pi - \varepsilon\theta) \setminus \pi$ and $\ddot{\pi} = (-\pi + x_1 + x_0) \setminus (-\pi + x_1 + x_0 - \varepsilon\theta)$; see Fig. 9.

In order to prove $g_{K,L}(x) \neq g_{K',L'}(x)$, we need to distinguish two cases, according to whether $[a_1, a_2]$ is an edge of K (or of K') or not. Note that $[a_1, a_2]$ is an edge of K if and only if $[b_1, b_2]$ is an edge of $-L'$. Indeed these conditions are equivalent respectively to $-\theta \in N(K, a_1)$ and to $-\theta \in N(-L', b_1)$, and these cones coincide by (5.4). Similar arguments prove that $[a_1, a_2]$ is an edge of K' if and only if $[b_1, b_2]$ is an edge of $-L$. We also observe that $[a_1, a_2]$ cannot be an edge of both K and K' , since otherwise $\Sigma = \partial K$, contradicting what has been proved in the lines preceding Claim 5.5.1.

Assume that $[a_1, a_2]$ is an edge of K . In this case $[b_1, b_2]$ is an edge of $-L'$, $\text{relint}[a_1, a_2] \subset K'$ and $\text{relint}[b_1, b_2] \subset -L$. We have $K \cap \dot{\pi} = \emptyset$, because $K \subset \pi$. When ε is small, the inclusion (5.25) implies $(L' + x) \cap \ddot{\pi} \subset \text{int } K'$, by continuity; see Fig. 9. Therefore

$$((K \cap (L + x)) \Delta (K' \cap (L' + x))) \cap \dot{\pi} = (L' + x) \cap \dot{\pi}.$$

This set is a rectangle of base $\|\dot{b}_2 - \dot{b}_1\|$ and height ε , up to triangles of edge-lengths proportional to ε . Its area is $\varepsilon \|\dot{b}_2 - \dot{b}_1\| + o(\varepsilon^2)$. Similar arguments prove that

$$((K \cap (L + x)) \Delta (K' \cap (L' + x))) \cap \ddot{\pi} = K \cap (L + x) \cap \ddot{\pi},$$

and that this set has area $\varepsilon \|-b_1 + x_0 - a_1\| + o(\varepsilon^2)$. Therefore we have

$$g_{K,L}(x) - g_{K',L'}(x) = \varepsilon (\|-b_1 + x_0 - a_1\| - \|\dot{b}_2 - \dot{b}_1\|) + o(\varepsilon^2),$$

which, in view of (5.28), contradicts $g_{K,L} = g_{K',L'}$. Assume that $[a_1, a_2]$ is neither an edge of K nor an edge of K' (and, as a consequence, $[b_1, b_2]$ is neither an edge of $-L$ nor an edge of $-L'$). In this case we have, for $\varepsilon > 0$ small,

$$((K \cap (L + x)) \Delta (K' \cap (L' + x))) \cap \dot{\pi} = \emptyset,$$

because $(L + x) \cap \dot{\pi} = (L' + x) \cap \dot{\pi}$ (by (5.29)) and both these sets are contained in K and K' (by (5.25)). Moreover, (5.29) implies $K \cap \ddot{\pi} = K' \cap \ddot{\pi}$ (because $\ddot{\pi} \subset \pi$ when ε is small), while (5.26) and (5.27) imply that $((L + x) \Delta (L' + x)) \cap \ddot{\pi}$ is contained in $B(-b_1 + x_0, \delta) \cup B(-b_2 + x_0, \delta)$, for a suitable $\delta = \delta(\varepsilon)$ positive which tends to 0 as ε tends to 0. If ε is sufficiently small, $B(-b_2 + x_0, \delta)$ does not intersect $K \cup K'$ and $B(-b_1 + x_0, \delta) \cap \ddot{\pi} \subset K, K'$. Thus we have

$$((K \cap (L + x)) \Delta (K' \cap (L' + x))) \cap \ddot{\pi} = ((L + x) \Delta (L' + x)) \cap B(-b_1 + x_0, \delta) \cap \ddot{\pi}.$$

Arguing as in the last part of the proof of Claim 5.5.2 proves $g_{K,L}(x) \neq g_{K',L'}(x)$. We omit the details.

Assume $\theta < \theta'$. The formulas (5.7), (5.8), (5.9), (5.10) and (5.21) imply that either $w_1 = \theta$ holds or $w_2 = \theta'$ holds. Assume $w_1 = \theta$, for instance. Let $\dot{a}_1, \dot{b}_1, \dot{a}_2, \dot{b}_2$ and π be defined as in case $\theta = \theta'$. Let us prove

$$(5.31) \quad \|a_2 - a_1\| < \|\dot{a}_2 - \dot{a}_1\|.$$

Assume (5.31) false and define $x_0 = a_2 + \dot{b}_2$. We have $[a_1, \dot{a}_1]_{\partial K} = [b_1, \dot{b}_1]_{\partial(-L)} + x_1$, $\dot{a}_1 = \dot{b}_1 + x_1$, $c_1 \in [\dot{a}_1, \dot{a}_2]_{\partial K} \cap ([\dot{b}_1, \dot{b}_2]_{\partial(-L)} + x_1)$ and $c_1 \neq \dot{a}_1$, because the arguments that prove these relations in the case $\theta = \theta'$ are valid also in this case. In particular $[\dot{b}_1, \dot{b}_2]$ is not a point. The condition $\theta < \theta'$ implies $-b_2 + x_0 \notin -\pi + x_1 + x_0$. Since $K, K' \subset -\pi + x_1 + x_0$, this implies $-b_2 + x_0 \notin K \cup K'$. Arguments similar to those used in the case $\theta = \theta'$ prove that $-\dot{b}_1 + x_0, -b_1 + x_0 \notin K \cup K'$. Therefore, when $x = x_0 - \varepsilon\theta$, with $\varepsilon > 0$ small, $(K \cap (L + x)) \Delta (K' \cap (L' + x))$ is contained in a neighbourhood of a_2 . Arguments similar to those in the last part of the proof

of Claim 5.5.2 prove $g_{K,L}(x) \neq g_{K',L'}(x)$. We omit the details. This contradiction proves (5.31). Arguments similar to those contained in the lines which follow (5.24) prove that (5.31) implies that U strictly contains an half-circle.

Let us prove $x_1 \neq x_2$ arguing by contradiction. If $x_1 = x_2$ then $\theta = \theta'$, by definition. Thus (5.22) and (5.23) hold and imply $\Sigma = \Omega + x_1$, contrary to the assumptions of Claim 5.5.4.

Summarising, U may coincide with an half-circle only when $\theta = \theta'$. When U is an half-circle, K is contained in the strip bounded by the line through a_1 orthogonal to u_1 and by the line through a_2 orthogonal to u_1 . Since $[a_1, a_2]$ is contained in this strip and it is parallel to $[a_1, a_2]$, equality holds in (5.24). Moreover the arcs $[a_1, a_1]_{\partial K}$ and $[a_2, a_2]_{\partial K}$ are line segments contained in the lines bounding the strip. The last part of the claim follows from these observations, (5.22) and (5.23). $\square$

Claim 5.5.5. *The arc Σ is a translate of Ω .*

Proof. Assume that Σ is not a translate of Ω . For $i = 1, 2$, let a_i, b_i and u_i be as in Claim 5.5.1, let $x_i = b_i - a_i$, let c_i be as in Claim 5.5.2 and let w_i be defined as in the lines preceding (5.21).

If $u \in (u_1, w_1)_{S^1}$, then both (2.3) and (2.4) hold, since $K_u, K'_u, (-L + x_1)_u$ and $(-L' + x_1)_u$ are all contained in the relative interior of $\Sigma \cap (\Omega + x_1)$, which is contained in $\partial K \cap \partial K' \cap (-\partial L + x_1) \cap (-\partial L' + x_1)$. Let $\tilde{U} \subset S^1$ be the maximal arc which contains $(u_1, w_1)_{S^1}$ and such that (2.4) holds for each $u \in \tilde{U}$. Let $\tilde{\Sigma}$ be the maximal arc of $\partial K \cap \partial(-L' + x_1)$ containing $\cup_{u \in (u_1, w_1)_{S^1}} K_u$ and let $\tilde{\Omega}$ be the maximal arc of $\partial K' \cap \partial(-L + x_1)$ containing $\cup_{u \in (u_1, w_1)_{S^1}} K'_u$. Since $u_1 \neq w_1$, by (5.7), $\tilde{U}$ is not a point and neither $\tilde{\Sigma}$ nor $\tilde{\Omega}$ are points or line segments. Clearly $\tilde{\Sigma}$ and $\tilde{\Omega}$ contain a_1 . Moreover, since ∂K and $\partial(-L' + x_1)$ bifurcate at c_1 (and the same is true for $\partial K'$ and $\partial(-L + x_1)$), c_1 is the upper endpoint of $\tilde{\Sigma}$ and of $\tilde{\Omega}$ and $\tilde{\Sigma} \neq \partial K$.

If $\tilde{\Sigma}$ is a translate of $\tilde{\Omega}$, then this translation is the identity, since $\tilde{\Sigma}$ and $\tilde{\Omega}$ have their upper endpoint in common. On the other hand, (5.4), with $j = 1$, implies that $\tilde{\Sigma}$ coincides with ∂K and $\tilde{\Omega}$ coincides with $\partial K'$ in a neighbourhood of a_1 . Since ∂K and $\partial K'$ bifurcate at a_1 , $\tilde{\Sigma}$ is not a translate of $\tilde{\Omega}$.

Results analogous to Claim 5.5.1 and 5.5.4 hold for $\tilde{U}, \tilde{\Sigma}$ and $\tilde{\Omega}$. In particular, $\tilde{U}$ contains an half-circle.

Let $\bar{U} \subset S^1$ be the maximal arc which contains $(w_2, u_2)_{S^1}$ and such that (2.4) holds for each $u \in \bar{U}$. Let $\bar{\Sigma}$ be the maximal arc of $\partial K \cap \partial(-L' + x_2)$ containing $\cup_{u \in (w_2, u_2)_{S^1}} K_u$ and let $\bar{\Omega}$ be the maximal arc of $\partial K' \cap \partial(-L + x_2)$ containing $\cup_{u \in (w_2, u_2)_{S^1}} K'_u$. Also $\bar{U}$ contains an half-circle.

For $i = 1, 2$, let $\tilde{u}_i$ and $\bar{u}_i \in S^1$ be such that $\text{cl } \tilde{U} = [\tilde{u}_1, \tilde{u}_2]_{S^1}$ and $\text{cl } \bar{U} = [\bar{u}_1, \bar{u}_2]_{S^1}$, and let $\tilde{a}_1 \in \partial K$ and $\bar{a}_2 \in \partial K$ be such that $\tilde{\Sigma} = [\tilde{a}_1, c_1]_{\partial K}$ and $\bar{\Sigma} = [c_2, \bar{a}_2]_{\partial K}$. Let us prove $\bar{u}_2 \leq \tilde{u}_1$. Assume $\bar{u}_2 > \tilde{u}_1$. In this case the sub-arcs $\tilde{\Sigma}$ and $\bar{\Sigma}$ of ∂K overlap and contain the arc $[\tilde{a}_1, \bar{a}_2]_{\partial K}$. The latter is not a point or a line segment, because arguing as we did in Claim 5.5.1 one can prove that $\tilde{U}$ contains a line segment orthogonal to $\tilde{u}_1$ containing $\tilde{a}_1$ and $\bar{U}$ contains a line segment orthogonal to $\bar{u}_2$ containing $\bar{a}_2$. The inclusions

$$[\tilde{a}_1, \bar{a}_2]_{\partial K} \subset \tilde{\Sigma} \subset \partial(-L' + x_1), \quad \text{and} \quad [\tilde{a}_1, \bar{a}_2]_{\partial K} \subset \bar{\Sigma} \subset \partial(-L' + x_2),$$

and the convexity of $\partial(-L')$ imply $x_1 = x_2$. This equality contradicts the assumption “ Σ is not a translate of Ω ”, as shown by Claim 5.5.4, and proves $\bar{u}_2 \leq \tilde{u}_1$. Similar arguments prove $\bar{u}_1 \geq \tilde{u}_2$. These inequalities imply that both $\tilde{U}$ and $\bar{U}$ are half-circles with $\tilde{U} \cup \bar{U} = S^1$.

A description analogous to that of Claim 5.5.4 applies to $\tilde{\Sigma}$ and $\tilde{\Omega}$ and also to $\bar{\Sigma}$ and $\bar{\Omega}$. This description easily implies that K , K' , $-L$ and $-L'$ are parallelograms with two edges orthogonal to $\tilde{u}_1$ and two edges orthogonal to v , for some $v \in S^1$ with $v \neq \tilde{u}_1$. It also implies $\lambda_1(K_v) = \lambda_1((-L')_v)$ and $\lambda_1(K'_v) = \lambda_1((-L)_v)$. It cannot be $\lambda_1(K_{\tilde{u}_1}) = \lambda_1((-L')_{\tilde{u}_1})$, because otherwise $\tilde{\Sigma} = \partial K$ and this contradicts what has been proved above. Thus, $\lambda_1(K_{\tilde{u}_1}) = \lambda_1(K'_{\tilde{u}_1})$ and $\lambda_1((-L)_{\tilde{u}_1}) = \lambda_1((-L')_{\tilde{u}_1})$, by the synisothesis of $(K, -L)$ and $(K', -L')$. Therefore, up to an affine transformation, (K, L) and (K', L') are trivial associates of $(\mathcal{K}_3, \mathcal{L}_3)$ and $(\mathcal{K}_4, \mathcal{L}_4)$ (with the defining parameter m equal to 0), respectively. This contradicts the assumptions of Lemma 5.5 and concludes its proof. $\square$

6. PROOF OF THEOREM 1.1

Proof. Proposition 4.2 implies that $(K', -L')$ is a pair of polygons synisothetic to $(K, -L)$. In particular, for each $u \in S^1$, either (2.3) or (2.4) holds. We assume that (K, L) and (K', L') are not trivial associates and prove that

$$(6.1) \quad K = -L + x \quad \text{and} \quad K' = -L' + x', \quad \text{for some } x, x' \in \mathbb{R}^2.$$

These identities, together with $K - L = K' - L'$ (which follows by (2.2)), imply $K = K' + (x - x')/2$ and $L = L' + (x - x')/2$, that is, they prove that (K, L) and (K', L') are trivial associates, concluding the proof.

In order to prove (6.1), let p be a vertex of K and q a vertex of $-L$ such that $\text{relint } N(K, p) \cap \text{relint } N(-L, q) \neq \emptyset$. We prove that

$$(6.2) \quad N(K, p) = N(-L, q).$$

Let $u_0 \in S^1 \cap \text{relint } N(K, p) \cap \text{relint } N(-L, q)$ and assume that (2.3) holds when $u = u_0$. This condition implies that there exist $y, y' \in \mathbb{R}^2$ such that K and $K' + y$ coincide in a neighbourhood of p , while $-L$ and $-(L' + y')$ coincide in a neighbourhood of q . Formulas (2.2) and (2.1) imply

$$p + q = K_{u_0} + (-L)_{u_0} = K'_{u_0} + (-L')_{u_0} = (p - y) + (q + y'),$$

that is $y = y'$. We apply Lemma 5.5 to (K, L) and $(K' + y, L' + y)$, with U chosen so that it contains $N(K, p) \cap N(-L, q)$. If Σ and Ω are defined as in the statement of Lemma 5.5, then they are not points nor line segments. This lemma implies that Σ is a translate of Ω , which yields (6.2). Similar arguments prove (6.2) when (2.4) replaces (2.3). In this case we apply Lemma 5.5 to (K, L) and $(-L' + y, -K' + y)$, where $y \in \mathbb{R}^2$ is chosen so that K and $-L' + y$ coincide in a neighbourhood of p .

What has been proved so far implies that to each edge E of K it corresponds an edge F of $-L$ with equal outer normal, and vice versa. To prove that K is a translate of $-L$ it suffices to show that

$$(6.3) \quad \lambda_1(E) = \lambda_1(F).$$

Let $E = [x_1, x_2]$ and $F = [y_1, y_2]$. We may label the vertices in such a way that $\text{int } N(K, x_i) \cap \text{int } N(-L, y_i) \neq \emptyset$, for each $i = 1, 2$. Let u_0 be the unit outer normal to K at E and assume that (2.3) holds when $u = u_0$. One proves, arguing as above, that there exists $y \in \mathbb{R}^2$ such that E is an edge of $K' + y$ with outer normal u_0 and F is an edge of $-(L' + y)$ with outer normal u_0 . We apply Lemma 5.5 to (K, L) and $(K' + y, L' + y)$, with U chosen so that it contains u_0 . What has been proved above implies that K and $K' + y$ coincide in a neighbourhood of E , that $-L$ and $-(L' + y)$ coincide in a neighbourhood of F , and that U is not a point. If Σ and Ω are defined as in the statement of Lemma 5.5, then they are not points nor line segments. This lemma implies that Σ is a translate of Ω , which yields (6.3). Similar arguments get the same conclusion when (2.4) substitutes (2.3), and similar arguments also prove that K' is a translate of $-L'$. $\square$

Proof of Corollary 1.2. First we prove the corollary assuming $z = 0$. Assume $g_{K,L}(x) = g_{K,L}(-x)$ for each $x \in \mathbb{R}^2$. This is equivalent to $g_{K,L}(x) = g_{L,K}(x)$ for each $x \in \mathbb{R}^2$, since $g_{L,K}(x) = g_{K,L}(-x)$. We claim that there exist no affine transformation $\mathcal{T}$ and no different indices i, j , with either $i, j \in \{1, 2\}$ or $i, j \in \{3, 4\}$, such that $(\mathcal{T}K, \mathcal{T}L)$ and $(\mathcal{T}L, \mathcal{T}K)$ are trivial associates of $(\mathcal{K}_i, \mathcal{L}_i)$ and $(\mathcal{K}_j, \mathcal{L}_j)$, respectively. Indeed, if this claim is false, then $(\mathcal{K}_i, \mathcal{L}_i)$ is a trivial associate of $(\mathcal{L}_j, \mathcal{K}_j)$, because being trivial associates is a transitive property. However, when $i \neq j$, $(\mathcal{K}_i, \mathcal{L}_i)$ is not a trivial associate of $(\mathcal{L}_j, \mathcal{K}_j)$, because $\mathcal{K}_i$ is not a translate of $-\mathcal{K}_j$ or of $\mathcal{L}_j$. This claim and Theorem 1.1 imply that (K, L) is a trivial associate of (L, K) . It is immediate to understand that this happens exactly when $K = -K + y$ and $L = -L + y$, for some $y \in \mathbb{R}^2$, (that is, $y/2$ is the center of K and of L) or when $K = L$. The converse implication follows from the identities $g_{K,L}(x) = g_{-K+y, -L+y}(-x) = g_{L,K}(-x)$, valid for any $x, y \in \mathbb{R}^2$.

The proof for $z \neq 0$ follows from the one for $z = 0$ applied to $g_{K,L+z}$, since $g_{K,L}(z+x) = g_{K,L+z}(x)$ and $g_{K,L}(z-x) = g_{K,L+z}(-x)$. $\square$

ACKNOWLEDGEMENTS. We are extremely grateful to G. Averkov and R. J. Gardner for reading large parts of this manuscript and suggesting many arguments that simplified and clarified some proofs. We also thank P. Mani-Levitska for giving us his unpublished note [Man01].

REFERENCES

- AP91. R. J. Adler and R. Pyke, *Problem 91–3*, Inst. Math. Statist. Bull. **20** (1991), 409.
- AP97. ———, *Scanning Brownian processes*, Adv. in Appl. Probab. **29** (1997), 295–326.
- AB07. G. Averkov and G. Bianchi, *Retrieving convex bodies from restricted covariogram functions*, Adv. in Appl. Probab. **39** (2007), 613–629.
- AB. ———, *Confirmation of Matheron’s conjecture on the covariogram of a planar convex body*, preprint, arXiv:0711.0572.
- BG07. M. Baake and U. Grimm, *Homometric model sets and window covariograms*, Zeitschrift für Kristallographie **222** (2007), 54–58.
- BBD. A. Benassi, G. Bianchi and G. D’Ercole, *Covariograms of non-convex sets*, in preparation.
- Bia05. G. Bianchi, *Matheron’s conjecture for the covariogram problem*, J. London Math. Soc. (2) **71** (2005), 203–220.
- Bia. ———, *The covariogram determines three-dimensional convex polytopes*, preprint.
- BSV00. G. Bianchi, F. Segala and A. Volčič, *The solution of the covariogram problem for plane C_+^2 bodies*, J. Differential Geom. **60** (2002), 177–198.
- CB03. A. Cabo and A. Baddeley, *Estimation of mean particle volume using the set covariance function*, Adv. in Appl. Probab. **35** (2003), 27–46.
- CJ94. A. Cabo and R. H. P. Janssen, *Cross-covariance functions characterise bounded closed regular sets*, CWI Technical Report, Amsterdam, 1994.
- GGZ05. R. J. Gardner, P. Gronchi and C. Zong, *Sums, projections, and sections of lattice sets, and the discrete covariogram*, Discrete Comput. Geom. **34** (2005), 391–409.
- Hor83. L. Hörmander, *The analysis of linear partial differential operators I*, Springer-Verlag, Berlin, 1983.
- Man01. P. Mani-Levitska, unpublished note, 2001.
- Mat75. G. Matheron, *Random sets and integral geometry*, Wiley, New York, 1975.
- Mat86. ———, *Le covariogramme géométrique des compacts convexes de $\mathbb{R}^2$* , Technical report 2/86, 54, Centre de Géostatistique, Ecole des Mines de Paris, 1986.
- MRG03. A. Mazzolo, B. Roesslinger and W. Gille, *Properties of chord length distributions of nonconvex bodies*, J. Math. Phys. **44** (2003), 6195–6208.
- Scm93. M. Schmitt, *On two inverse problems in mathematical morphology*, Mathematical Morphology in Image Processing, Dekker, New York, 1993, pp. 151–169.
- Sch93. R. Schneider, *Convex bodies: the Brunn-Minkowski theory*, Cambridge University Press, Cambridge, 1993.
- Ser84. J. Serra, *Image analysis and mathematical morphology*. Academic Press, London, 1984.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, FIRENZE,
ITALY I-50134

E-mail address: `gabriele.bianchi@unifi.it`